

Title: Glucoamylase variants

CROSS-REFERENCE TO RELATED APPLICATIONS

5 This application is a division of U.S. patent application 09/351,814, filed on July 12, 2001, and claims priority under 35 U.S.C. 119 of Danish application nos. PA 1998 00937 and PA 19980167 filed on July 15, 1998 and December 17, 1998, respectively, and U.S provisional application nos. 60/093,528 and 60/115,545 filed on July 21, 1998 and January 12, 1999, respectively, the contents of which are fully incorporated herein
10 by reference.

FIELD OF THE INVENTION

15 The present invention relates to novel glucoamylase variants (mutants) of parent AMG, in particular with improved thermal stability and/or increased specific activity suitable for, e.g., starch conversion, e.g., for producing glucose from starch. More specifically, the present invention relates to glucoamylase enzyme variants and the use of such variant enzymes.

BACKGROUND OF THE INVENTION

20 Glucoamylase (1,4- α -D-glucan glucohydrolase, EC 3.2.1.3) is an enzyme which catalyzes the release of D-glucose from the non-reducing ends of starch or related oligo- and polysaccharide molecules. Glucoamylases are produced by several filamentous fungi and yeasts, with those from *Aspergillus* being commercially most important.

25 Commercially, the glucoamylase enzyme is used to convert corn starch which is already partially hydrolyzed by an α -amylase to glucose. The glucose is further converted by glucose isomerase to a mixture composed almost equally of glucose and fructose. This mixture, or the mixture further enriched with fructose, is the commonly used high fructose corn syrup commercialized throughout the world. This syrup is the world's largest tonnage product produced by an enzymatic process. The three enzymes involved in the
30 conversion of starch to fructose are among the most important industrial enzymes produced.

One of the main problems that exist with regard to the commercial use of glucoamylase in the production of high fructose corn syrup is the relatively low thermal stability of glucoamylase. Glucoamylase is not as thermally stable as α -amylase or
35 glucose isomerase and it is most active and stable at lower pH's than either α -amylase or

glucose isomerase. Accordingly, it must be used in a separate vessel at a lower temperature and pH.

Glucoamylase from *Aspergillus niger* has a catalytic (aa 1-440) and a starch binding domain (aa 509-616) separated by a long and highly O-glycosylated linker (Svensson et al. (1983), *Carlsberg Res. Commun.* **48**, 529-544, 1983 and (1986), *Eur. J. Biochem.* **154**, 497-502). The catalytic domain (aa 1-471) of glucoamylase from *A. awamori* var. X100 adopt an $(\alpha/\alpha)_6$ -fold in which six conserved $\alpha \rightarrow \alpha$ loop segments connect the outer and inner barrels (Aleshin et al. (1992), *J. Biol.Chem.* **267**, 19291-19298). Crystal structures of glucoamylase in complex with 1-deoxynojirimycin (Harris et al. (1993), *Biochemistry*, **32**, 1618-1626) and the pseudotetrasaccharide inhibitors acarbose and D-gluco-dihydroacarbose (Aleshin et al. (1996), *Biochemistry* **35**, 8319-8328) furthermore are compatible with glutamic acids 179 and 400 acting as general acid and base, respectively. The crucial role of these residues during catalysis have also been studied using protein engineering (Sierks et al. (1990), *Protein Engng.* **3**, 193-198; Frandsen et al. (1994), *Biochemistry*, **33**, 13808-13816). Glucoamylase-carbohydrate interactions at four glycosyl residue binding subsites, -1, +1, +2, and +3 are highlighted in glucoamylase-complex structures (Aleshin et al. (1996), *Biochemistry* **35**, 8319-8328) and residues important for binding and catalysis have been extensively investigated using site-directed mutants coupled with kinetic analysis (Sierks et al. (1989), *Protein Engng.* **2**, 621-625; Sierks et al. (1990), *Protein Engng.* **3**, 193-198; Berland et al. (1995), *Biochemistry*, **34**, 10153-10161; Frandsen et al. (1995), *Biochemistry*, **34**, 10162-10169).

Different substitutions in *A. niger* glucoamylase to enhance the thermal stability have been described: i) substitution of α -helical glycines: G137A and G139A (Chen et al. (1996), *Prot. Engng.* **9**, 499-505); ii) elimination of the fragile Asp-X peptide bonds, D257E and D293E/Q (Chen et al. (1995), *Prot. Engng.* **8**, 575-582); prevention of deamidation in N182 (Chen et al. (1994), *Biochem. J.* **301**, 275-281); iv) engineering of additional disulphide bond, A246C (Fierobe et al. (1996), *Biochemistry*, **35**, 8698-8704; and v) introduction of Pro residues in position A435 and S436 (Li et al. (1997), *Protein Engng.* **10**, 1199-1204. Furthermore Clark Ford presented a paper on Oct 17, 1997, ENZYME ENGINEERING 14, Beijing/China Oct 12-17, 97, Abstract number: Abstract book p. 0-61. The abstract suggests mutations in positions G137A, N20C/A27C, and S30P in a (not disclosed) *Aspergillus awamori* glucoamylase to improve the thermal stability.

Additional information concerning glucoamylase can be found on an Internet homepage

(<http://www.public.iastate.edu/~pedro/glase/glase.html>) "Glucoamylase WWW page" (Last changed 97/10/08) by Pedro M. Coutinho discloses informations concerning glucoamylases, including glucoamylases derivable from *Aspergillus* strains. Chemical and site-directed modifications in the *Aspergillus niger* glucoamylase are listed.

5

BRIEF DISCLOSURE OF THE INVENTION

The object of the present invention is to provide improved glucoamylase variants with improved thermostability and/or increased specific activity suitable for use in, e.g., the saccharification step in starch conversion processes.

10 The term "a glucoamylase variant with improved thermostability" means in the context of the present invention a glucoamylase variant which has a higher $T_{1/2}$ (half-time) than the corresponding parent glucoamylase. The determination of $T_{1/2}$ (Method I and Method II) is described below in the "Materials & Methods" section.

15 The term "a glucoamylase variant with increased specific activity" means in the context of the present invention a glucoamylase variant with increased specific activity towards the α -1,4 linkages in the saccharide in question. The specific activity is determined as k_{cat} or AGU/mg (measured as described below in the "Materials & Methods" section). An increased specific activity means that the k_{cat} or AGU/mg values are higher when compared to the k_{cat} or AGU/mg values, respectively, of the
20 corresponding parent glucoamylase.

The inventors of the present invention have provided a number of improved variants of a parent glucoamylase with improved thermostability and/or increased specific activity in comparison to the parent corresponding enzyme. The improved thermal stability is obtained by substituting selected positions in a parent glucoamylase. This will be
25 described in details below.

Nomenclature

In the present description and claims, the conventional one-letter and three-letter codes for amino acid residues are used. For ease of reference, glucoamylase variants of
30 the invention are described by use of the following nomenclature:

Original amino acid(s):position(s):substituted amino acid(s)

According to this nomenclature, for instance the substitution of alanine for asparagine in position 30 is shown as:

Ala30Asn or A30N

35 a deletion of alanine in the same position is shown as:

Ala30* or A30*

and insertion of an additional amino acid residue, such as lysine, is shown as:

Ala30AlaLys or A30AK

A deletion of a consecutive stretch of amino acid residues, such as amino acid residues 30-33, is indicated as (30-33)* or Δ (A30-N33).

5 Where a specific glucoamylase contains a "deletion" in comparison with other glucoamylases and an insertion is made in such a position this is indicated as:

*36Asp or *36D

for insertion of an aspartic acid in position 36

Multiple mutations are separated by plus signs, *i.e.*:

10 Ala30Asp + Glu34Ser or A30N+E34S

representing mutations in positions 30 and 34 substituting alanine and glutamic acid for asparagine and serine, respectively. Multiple mutation may also be separated as follows, *i.e.*, meaning the same as the plus sign:

Ala30Asp/Glu34Ser or A30N/E34S

15 When one or more alternative amino acid residues may be inserted in a given position it is indicated as

A30N,E or A30N/E, or A30N or A30E

Furthermore, when a position suitable for modification is identified herein without any specific modification being suggested, it is to be understood that any amino acid residue
20 may be substituted for the amino acid residue present in the position. Thus, for instance, when a modification of an alanine in position 30 is mentioned, but not specified, it is to be understood that the alanine may be deleted or substituted for any other amino acid, *i.e.*, any one of:

R,N,D,A,C,Q,E,G,H,I,L,K,M,F,P,S,T,W,Y,V.

25

BRIEF DESCRIPTION OF THE DRAWING

Figure 1 shows the plasmid pCAMG91 containing the *Aspergillus niger* G1 glucoamylase gene.

30 DETAILED DISCLOSURE OF THE INVENTION

A goal of the work underlying the present invention was to improve the thermal stability and/or increase the specific activity of particular glucoamylases which are obtainable from fungal organisms, in particular strains of the *Aspergillus* genus and which themselves had been selected on the basis of their suitable properties in starch
35 conversion or alcohol fermentation.

Identifying positions and/or regions to be mutated to obtain Improved Thermostability and/or Increased Specific Activity

Molecular dynamics (MD) simulations indicate the mobility of the amino acids in the protein structure (see McCammon, JA and Harvey, SC., (1987), "Dynamics of proteins and nucleic acids". Cambridge University Press.). Such protein dynamics are often compared to the crystallographic B-factors (see Stout, GH and Jensen, LH, (1989), "X-ray structure determination", Wiley). By running the MD simulation at different protonation states of the titrate able residues, the pH related mobility of residues are simulated. Regions having the highest mobility or flexibility (here isotropic fluctuations) are selected for random mutagenesis. It is here understood that the high mobility found in certain areas of the protein, can be thermally improved by substituting residues in these residues. The substitutions are directed against residues that will change the dynamic behaviour of the residues to e.g. bigger side-chains and/or residues which have capability of forming improved contacts to residues in the near environment. The AMG from *Aspergillus niger* was used for the MD simulation. How to carry out MD simulation is described in the Materials & Methods" section below.)

Regions found by Molecular dynamics (MD) simulations to be suitable for mutation when wanting to obtain improved thermal stability and/or increased specific activity are the following:

Region: 1-18,
Region: 19-35,
Region: 73-80,
Region: 200-212,
Region: 234-246,
Region: 334-341,
Region: 353-374,
Region: 407-411,
Region: 445-470,

Regions found to be of interest for increasing the specific activity and/or improved thermostability are the regions in proximity to the active site. Regions positioned in between the α -helixes, and which may include positions on each side of the N- and C-terminal of the α -helixes, at the substrate binding site is of importance for the activity of the enzyme. These regions constitute the following regions:

Region: 40-62,
Region: 93-127,
Region: 170-184,

Region: 234-246,
Region: 287-319,
Region: 388-414.

Rhizopus, *Talaromyces*, such as *Talaromyces emersonii* (disclosed in WO 99/28448), and *Thielavia* have high specific activity towards maltodextrins, including maltose and maltohepatose. Therefore, regions being of special interest regarding (transferring) increased specific activity are:

Region: 200-212,
Region: 287-300,
Region: 305-319.

The present inventors have found that it is in fact possible to improve the thermal stability and/or to increase the specific activity of a parent glucoamylase by modification of one or more amino acid residues of the amino acid sequence of the parent glucoamylase. The present invention is based on this finding.

Accordingly, in a first aspect the present invention relates to an improved variant of a parent glucoamylase comprising one or more mutations in the regions and positions described further below.

Parent Glucoamylases

Parent glucoamylase contemplated according to the present invention include fungal glucoamylases, in particular fungal glucoamylases obtainable from an *Aspergillus* strain, such as an *Aspergillus niger* or *Aspergillus awamori* glucoamylases and variants or mutants thereof, homologous glucoamylases, and further glucoamylases being structurally and/or functionally similar to SEQ ID NO: 2. Specifically contemplated are the *Aspergillus niger* glucoamylases G1 and G2 disclosed in Boel et al. (1984), "Glucoamylases G1 and G2 from *Aspergillus niger* are synthesized from two different but closely related mRNAs", EMBO J. 3 (5), p. 1097-1102,. The G2 glucoamylase is disclosed in SEQ ID NO: 2. The G1 glucoamylase is disclosed in SEQ ID NO: 13 . Another AMG backbone contemplated is *Talaromyces emersonii* , especially *Talaromyces emersonii* DSM disclosed in WO 99/28448 (Novo Nordisk).

Commercially Available Parent Glucoamylases

Commercially available parent glucoamylases include AMG from Novo Nordisk, and also glucoamylase from the companies Genencor, Inc. USA, and Gist-Brocades, Delft, The Netherlands.

Glucoamylase Variants

In the first aspect the invention relates to a variant of a parent glucoamylase comprising one or more mutation(s) in the following position(s) or region(s) in the amino acid sequence shown in NO: 2:

- 5 Region: 1-18,
Region: 19-35,
Region: 40-62,
Region: 73-80,
Region: 93-127,
10 Region: 170-184,
Region: 200-212,
Region: 234-246,
Region: 287-319,
Region: 334-341,
15 Region: 353-374,
Region: 388-414,
Region: 445-470,

and/or in a corresponding position or region in a homologous glucoamylase which displays at least 60% homology with the amino acid sequences shown in SEQ ID NO: 2, with the exception of the following substitutions: N20C, A27C, S30P, Y48W, Y50F, W52F, R54K/L, D55G/V, G57A, K108R, D112Y, Y116A/W, S119C/W/E/G/Y/P, W120H/L/F/Y, G121T/A, R122Y, P123G, Q124H, R125K, W170F, N171S, Q172N, T173G, G174C, Y175F, D176N/E, L177H/D, W178R/D, E179Q/D, E180D/Q, V181D/A/T, N182A/D/Q/Y/S, G183K, S184H, W212F, R241K, A246C, D293E/Q, A302V, R305K, 25 Y306F, D309N/E, Y312W, W317F, E389D/Q, H391W, A392D, A393P, N395Q, G396S, E400Q/C, Q401E, G407D, E408P, L410F, S411A/G/C/H/D, S460P

In an embodiment the region mutated is the Region: 1-18.

Specific preferred positions contemplated include one or more of:

1,2,3,4,5,6,7,8,9,10,11,12,13,14,15,16,17,18.

- 30 Specific mutations include one or more of: A1V, T2P/Q/R/H/M/E/K, L3N, N9A, A11E/P, I18V.

Preferred combinations of mutations include one or more of:

- A1V+L66R+Y402F+N427S+S486G,
T2K+S30P+N427M+S44G+V470M,
35 T2E+T379A+S386K+A393R,
T2Q+A11P+S394R,

T2R+L66V+S394P+Y402F
T2M+N9A+T390R+D406N+L410R,
T2R+S386R+A393R,
A11P+T2Q+S394R,
5 A11E+E408R,
I18V+T51S+S56A+V59T+L60A.

In an embodiment the region mutated is the Region: 19-35.

Preferred sub-regions include one or more of: 21-26, 31-35.

Specific preferred positions contemplated include one or more of: 19, 21, 22, 23, 24, 25,
10 26, 28, 29, 31, 32, 33, 34, 35.

Specific mutations include one or more of: L19N, N20T, G23A, A24S/T, D25S/T/R,
G26A, A27S/T, W28R/Y, S30T/N, G31A, A32V, D33R/K/H, S34N.

In an embodiment the region mutated is the Region: 40-62.

Preferred sub-regions include one or more of: 40-47, 58-62.

15 Specific preferred positions contemplated include one or more of:
40,41,42,43,44,45,46,47,49,51,53,56,58,59,60,61,62.

Specific mutations include one or more of:

S40C/A/G,

T43R,

20 T51S/D,

T53D,

S56A/C,

V59T/A,

L60A.

25 Preferred combinations of mutations include one or more of:

T51S+S56A+V59T+L60A+I18V

V59A+A393R+T490A+PLASD(N-terminal extension).

In an embodiment the region mutated is the Region: 73-80.

Specific preferred positions contemplated include one or more of:
30 73,74,75,76,77,78,79,80.

Specific mutations include one or more of:

S73P/D/T/N/Q/E,

L74I/D,

L75A/R/N/D/C/Q/E/G/H/I/K/M/P/S/T/V, preferred are I/N/D,

35 S76A/R/N/D/C/Q/E/G/H/I/L/K/F/M/P/S/T/W/Y/V, preferred T/A)

T77V/T,

I78V,
E79A/R/N/D/C/Q/G/H/I/L/K/F/M/P/S/T/Y/V, preferred are Q/R/K)
N80A/R/D/C/Q/E/G/H/I/L/K/F/M/P/S/T/W/Y/V, preferred are
H/D/E/R/K/T/S/Y.

- 5 In an embodiment the region mutated is the Region: 93-127.
In an additional embodiment the sub-region is: Region: 93-124.
Preferred sub-regions include one or more of: 93-107, 109-111, 113-115.

Specific preferred positions contemplated include one or more of:
93,94,95,96,97,98,99,100,101,102,103,104,105,106,109,110,111,113,114,115,117,118,1
10 26,127.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

Specific mutations include one or more of: N93T,
P94V,
S95N,
D97S,
15 L98S/P,
S100T/D,
A102S/*,
P107M/L/A/G/S/T/V/I,
N110T,
20 V111P,
D112N,
E113M/A,
T114S,
A115Q/A,
25 Y116F,
S119A/R/N/D/Q/H/I/L/K/F/M/T/V, preferred is A,
R122A/N/D/C/Q/E/G/H/I/L/K/F/M/P/S/T/W/V,
G127A.

Preferred combinations of mutations include one or more of: S119P+G447S,

- 30 S119P+Y402F,
S119P+A393R,
S119P+I189T+223F+F227Y+Y402F
S119P+T416H+Y402F+Y312Q.

In an embodiment the region mutated is the Region: 170-184.

- 35 Specific mutations include one or more of:
N171R/K/Q/E/W/F/Y,

Q172A/R/N/D/C/E/G/H/I/L/K/F/M/P/S/T/W/Y/V,
T173K/R/S/N/Q,
G174A,S,
Y175N/Q/D/E,

5 D176L

L177I

E180N/M

V181I/T

N182R/C/E/G/H/I/L/K/M/P/T/W/Y/V.

10 G183A

S184D/N/E/Q/L/I/T/R/K.

In an embodiment the region mutated is the Region: 200-212.

Preferred sub-regions include: 200-211.

Specific preferred positions contemplated include one or more of:

15 200,201,202,203,204,205,206,207,208,209,210,211.

Specific mutations include one or more of: A201D,, F202L, A203L , T204K, A205R/S,
V206L/N, G207N, S208H/T/D, S209T, S211P, W212N/A/T.

In an embodiment the region mutated is the Region: 234-246.

Preferred sub-regions include one or more of: 234-240, 242-245.

20 Specific preferred positions contemplated include one or more of:

234,235,236,237,238,239,240,242,243,244,245.

Specific mutations include one or more of:

L234A/R/N/D/C/Q/E/G/H/I/K/M/F/P/S/T/W/Y/V.

A235S

25 F237Y/H/N/D.

D238T/S.

S239A/R/N/D/C/G/H/I/L/F/P/S/T/Y/V.

S240G

S242S/P/T/A/Y/H/N/D.

30 G243S/P/T/A/Y/H/N/D.

K244R,

A246T.

Preferred combinations of mutations include one or more of: A246T+T72I.

In an embodiment the region mutated is the Region: 287-319.

35 Preferred sub-regions include one or more of: 287-292, 294-301, 313-316.

Specific preferred positions contemplated include one or more of:
287,288,289,290,291,292,294,295,296,297,298,299,300,301,
303,307,308,310,311,313,314,315,316,318,319.

Specific mutations include one or more of:

- 5 S287A/R/N/D/C/Q/E/G/H/I/L/K/M/T/V,
I288L/N/Q,
Y289F,
T290A/R/N/D/C/Q/E/G/H/I/L/K/M/P/S/V,
L291I/D/N,
10 N292D,
D293A/R/N/C/Q/E/G/H/I/L/K/M/S/T/V,
G294A/R/N/D/C/Q/E/H/I/L/K/M/P/S/T/V,
L295A/R/N/D/C/Q/E/G/H/K/M/S/T/V,
S296A/R/N/D/C/Q/E/G/H/I/L/K/M/T/V,
15 D297A/R/N/C/Q/E/G/H/I/L/K/M/P/S/T/V,
S298A/R/N/D/C/Q/E/G/H/I/L/K/F/M/T/V,
E299A/R/N/D/C/Q/G/H/I/L/K/M/S/T/V.
V301T/I,
A302R/N/D/C/Q/E/G/H/I/L/K/F/M/P/S/T/W/Y/V/, preferred S, V303T/I,
20 G304A,
R305A/N/D/C/Q/E/G/H/I/L/F/M/P/S/T/W/Y/V,
Y306A/R/N/D/C/Q/E/G/H/I/L/K/M/P/S/T/W/V.
E308A/R/N/D/C/Q/G/H/I/L/K/M/P/S/T/V, preferred Q,
D309L,
25 T310V/S,
Y311N,
Y312Q/N,
N313T/S/G
N315Q/E/R,
30 F318A/R/N/D/C/Q/E/G/H/I/L/K/M/P/S/T/W/Y/V, preferred is Y.
Preferred combinations of mutations include one or more of:
Y312Q+S119P+T416H+Y402F,
Y312Q+S119P+Y402F+T416H+S411V,
Y312Q+T416H
35 N313G+F318Y.

In an embodiment the region mutated is the Region: 334-341.

Specific preferred positions contemplated include one or more of:

334,335,336,337,338,339,340,341.

Specific mutations include one or more of:

D336A/R/N/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V.

5 K337A/R/N/D/C/Q/E/G/H/I/L/M/F/P/S/T/W/Y/V.

Q338A/R/N/D/C/G/H/I/L/F/P/S/T/Y/V.

G339S/P/T/A.

S340I/T/N/V/A/D/G.

L341F/L/I/V.

10 Preferred combinations of mutations include one or more of:

S340G+D357S+T360V+S386P.

In an embodiment the region mutated is the Region: 353-374.

Specific preferred positions contemplated include one or more of:

353,354,355,356,357,358,359,360,361,362,363,364,365,366,367,368,369,370,371,372,3

15 73,374.

Specific mutations include one or more of:

A353D/S,

S356P/N/D,

D357S,

20 A359S,

T360V,

G361S/P/T/A,

T362R,

S364A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/T/W/Y/V,

25 S365A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/T/W/Y/V,

S366T,

S368P/T/A,

T369A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/T/W/Y/V,

S371Y/H/N/D,

30 S372F/Y/C/L/P/H/R/I/T/N/S/V/A/D/G.

Preferred combinations of mutations include one or more of:

S356P+S366T,

D357S+T360V+S371H.

D357S+T360V+S386P+S340G.

35 In an embodiment the region mutated is the Region: 388-414.

Preferred sub-regions include one or more of: 397-399,402-406, 412-414.

Specific preferred positions contemplated include one or more of:
388,389,390,394,397,398,399,402,403,404,405,406,409,412,413,414.

Specific mutations include one or more of: T390R,

A393R,

5 S394P/R,

M398L,

S399A/R/N/D/C/Q/E/G/H/I/L/K/F/M/P/T/W/Y/V, preferred are T/Q/C,

Y402A/R/N/D/C/Q/E/G/H/I/L/K/F/M/P/S/T/V, preferred is F,

D403S,

10 S405T,

D406N,

E408C/R,

A409R/N/D/C/Q/E/G/H/I/L/K/F/M/P/S/T/W/Y/V, preferred is P,

L410R/I,

15 S411R/N/Q/E/I/L/K/F/M/P/T/W/Y/V, preferred is V,

A412C,

R413A/N/D/C/Q/E/G/H/I/L/K/F/M/P/S/T/W/Y/V.

D414A.

Preferred combinations of mutations include one or more of:

20 A393R+T490A+V59A+PLASD(N-terminal extension)

S394R+T2Q+A11P,

Y402F+S411V,

Y402F+S411V+S119P

Y402F+S411V+S119P+A393R,

25 Y402F+Y312Q+S119P+T416H,

E408R+S386N,

E408R+A425T+S465P+T494A,

L410R+A393R,

Y402F+Y312Q+S119P+T416H+S411V+A393R.

30 In an embodiment the region mutated is the Region: 445-470.

Preferred sub-regions include one or more of: 445-459, 461-470.

Specific preferred positions contemplated include one or more of:

445,446,447,448,449,450,451,452,453,454,455,456,457,458,459,461,

462,463,464,465,466,467,467,468,469,470.

35 Specific mutations include one or more of: G447S,G456C/P,S465P.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30

Specific variants include variants having one or more of the following substitutions:

A1V, T2E/P/Q/R/H/M, L3P/N, N9A, A11P/E, I18V, L19N, N20T, G23A, A24S/T, D25S/T/R, G26A, A27S/T, W28R/Y, S30T/N, G31A, A32V, D33R/K/H, S34N, S40C, T43R, T51D/S, T53D, S56A/C, V59T/A, L60A, N93T, P94V, S95N, D97S, L98P/S, S100T/D, A102S/*, N110T, V111P, D112N, E113M/A, T114S, A115Q/G, Y116F, S119A, G127A, N182E, A201D, F202L, A203L, T204K, A205R/S, V206L/N, G207N, S208H/T/D, S209T, S211P, W212N/A/T, A246T Y312Q, N313T/S/G, A353D/S, S356P/N/D, D357S, A359S, T360V, G361S/P/T/A, T362R, S364A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/T/W/Y/V, S365A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/T/W/Y/V, S366T, S368P/T/A, T369A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/T/W/Y/V, S371Y/H/N/D, S372F/Y/C/L/P/H/R/I/T/N/S/V/A/D/G, T390R, A393R, S394R/P, M398L, S399C/Q/T, Y402F, D403S, S405T, D406N, E408C/R, L410I/R, S411V, A412C, D414A, G447S, S465P.

Improved thermal stability

In a second aspect the invention relates to a variant of a parent glucoamylase with improved thermal stability, in particular in the range from 40-80°C, preferably 60-80°C, and preferably at pH 4-5, said variant comprising one or more mutation(s) in the following position(s) or region(s) in the amino acid sequence shown in NO: 2:

Region: 1-18,

Region: 19-35,

Region: 73-80,

Region: 93-127,

Region: 170-184,

Region: 200-212,

Region: 234-246,

Region: 287-319,

Region: 334-341,

Region: 353-374,

Region: 388-414,

Region: 445-470,

and/or in a corresponding position or region in a homologous glucoamylase which displays at least 60% homology with the amino acid sequences shown in SEQ ID NO: 2, with the exception of the following substitutions: N20C, A27C, S30P, A246C.

As substrate binding may improve the stability region 93-127, Region: 170-184, Region: 305-319 are also contemplated for thermostabilization according to the present invention.

In an embodiment the region mutated is the Region: 1-18.

5 Specific preferred positions contemplated include one or more of:

1,2,3,4,5,6,7,8,9,10,11,12,13,14,15,16,17,18.

Specific mutations include one or more of: A1V, T2P/Q/R/H/M/E, N9A, A11E/P.

Preferred combinations of mutations include one or more of:

A1V+L66R+Y402F+N427S+S486G,

10 T2K+S30P+N427M+S44G+V470M,

T2E+T379A+S386K+A393R,

T2R+S386R+A393R,

A11P+T2Q+S394R,

T2Q+A11P+S394R,

15 T2R+L66V+S394P+Y402F,

T2M+N9A+T390R+D406N+L410R,

T2R+S386R+A393R,

A11E+E408R.

In an embodiment the region mutated is the Region: 19-35.

20 Preferred sub-regions include one or more of: 21-26, 31-35.

Specific preferred positions contemplated include one or more of:

19,21,22,23,24,25,26,28,29,31,32,33,34,35.

Specific mutations include one or more of: L19N, N20T, G23A, A24S/T, D25S/T/R, G26A, A27S/T, W28R/Y, S30T/N, G31A, A32V, D33R/K/H, S34N.

25 In an embodiment the region mutated is the Region: 73-80.

Specific preferred positions contemplated include one or more of:

73,74,75,76,77,78,79,80.

Specific mutations include one or more of:

S73P/D/T/N/Q/E,

30 L74I/D,

L75A/R/N/D/C/Q/E/G/H/I/K/M/P/S/T/V, preferred are I/N/D,

S76A/R/N/D/C/Q/E/G/H/I/L/K/F/M/P/S/T/W/Y/V, preferred T/A)

T77V/T,

I78V,

35 E79A/R/N/D/C/Q/G/H/I/L/K/F/M/P/S/T/Y/V, preferred are Q/R/K)

N80A/R/D/C/Q/E/G/H/I/L/K/F/M/P/S/T/W/Y/V, preferred are

H/D/E/R/K/T/S/Y.

In an embodiment the region mutated is the Region: 93-127.

In an additional embodiment the sub-region is: Region: 93-124

Preferred sub-regions include one or more of: 93-107, 109-111, 113-115.

5 Specific preferred positions contemplated include one or more of:

93,94,95,96,97,98,99,100,101,102,103,104,105,106,109,110,111,113,114,115,117,118,126,127.

Preferred mutations include one or more of:

P107M/L/A/G/S/T/V/I,

10 S119A/R/N/D/Q/H/I/L/K/F/M/T/V, preferred is A,

R122A/N/D/C/Q/E/G/H/I/L/K/F/M/P/S/T/W/V,

Preferred combinations of mutations include one or more of: S119P+G447S.

S119P+A393R.

In an embodiment the region mutated is the Region: 170-184.

15 Specific mutations include one or more of:

N171R/K/Q/E/W/F/Y,

Q172A/R/N/D/C/E/G/H/I/L/K/F/M/P/S/T/W/Y/V,

T173K/R/S/N/Q,

G174A,S,

20 Y175N/Q/D/E,

D176L

L177I

E180N/M

V181I/T

25 N182R/C/E/G/H/I/L/K/M/P/T/W/Y/V.

G183A

S184D/N/E/Q/L/I/T/R/K.

In an embodiment the region mutated is the Region: 200-212.

Preferred sub-regions include: 200-211.

30 Specific preferred positions contemplated include one or more of:

200,201,202,203,204,205,206,207,208,209,210,211.

Specific mutations include one or more of: A203L, S211P.

In an embodiment the region mutated is the Region: 234-246.

Preferred sub-regions include one or more of: 234-240, 242-245.

35 Specific preferred positions contemplated include one or more of:

234,235,236,237,238,239,240,242,243,244,245.

Specific mutations include one or more of:

L234A/R/N/D/C/Q/E/G/H/I/K/M/F/P/S/T/W/Y/V,

A235S,

F237Y/H/N/D,

5 D238T/S,

S239A/R/N/D/C/G/H/I/L/F/P/S/T/Y/V,

S240G,

S242S/P/T/A/Y/H/N/D,

G243S/P/T/A/Y/H/N/D,

10 K244R,

A246T.

Preferred combinations of mutations include one or more of: A246T+T72I.

In an embodiment the region mutated is the Region: 287-319.

Preferred sub-regions include one or more of: 287-292, 294-301, 313-316.

15 In an additional embodiment the sub-region include: 305-319

Specific preferred positions contemplated include one or more of:

287,288,289,290,291,292,294,295,296,297,298,299,300,301,302,

303,307,308,310,311,313,314,315,316,318,319.

Specific mutations include one or more of:

20 Y312Q,

F318A/R/N/D/C/Q/E/G/H/I/L/K/M/P/S/T/W/Y/V, preferred is Y.

Preferred combinations of mutations include one or more of:

N313G+F318Y,

Y302Q+S119P+T416H+Y402F.

25 In an embodiment the region mutated is the Region: 334-341.

Specific preferred positions contemplated include one or more of:

334,335,336,337,338,339,340,341.

Specific mutations include one or more of:

D336A/R/N/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V.

30 K337A/R/N/D/C/Q/E/G/H/I/L/M/F/P/S/T/W/Y/V.

Q338A/R/N/D/C/G/H/I/L/F/P/S/T/Y/V.

G339S/P/T/A.

S340I/T/N/V/A/D/G.

L341F/L/I/V.

35 Preferred combinations of mutations include one or more of:

S340G+D357S+T360V+S386P.

In an embodiment the region mutated is the Region: 353-374.

Specific preferred positions contemplated include one or more of:

353,354,355,356,357,358,359,360,361,362,363,364,365,366,367,368,
369,370,371,372,373,374.

5 Specific mutations include one or more of:

A353D/S,

S356P/N/D,

D357S,

A359S,

10 T360V,

G361S/P/T/A,

T362R,

S364A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/T/W/Y/V,

S365A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/T/W/Y/V,

15 S366T,

S368P/T/A,

T369A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/T/W/Y/V,

S371Y/H/N/D,

S372F/Y/C/L/P/H/R/I/T/N/S/V/A/D/G.

20 Preferred combinations of mutations include one or more of:

S356P+S366T,

D357S+T360V+S371H

D357S+T360V+S386P+S340G.

In an embodiment the region mutated is the Region: 388-414.

25 Preferred sub-regions include one or more of: 397-399,402-406, 412-414.

In an additional embodiment the sub-region is: 407-411

Specific preferred positions contemplated include one or more of:

388,389,390,394, 397,398,399,402,403,404,405,406,409,412,413,414.

Specific mutations include one or more of: T390R,

30 A393R,

S394P/R,

S399A/R/N/D/C/Q/E/G/H/I/L/K/F/M/P/T/W/Y/V, preferred are T/Q/C,

Y402A/R/N/D/C/Q/E/G/H/I/L/K/F/M/P/S/T/V, preferred is F,

D403S,

35 S405T,

D406N,

E408C/R,
Q409A/R/N/D/C/E/G/H/I/L/K/F/M/P/S/T/W/Y/V, preferred is P,
L410I/R,
S411V,
5 A412C,
R413A/N/D/C/Q/E/G/H/I/L/K/F/M/P/S/T/W/Y/V.
D414A.

Preferred combinations of mutations include one or more of:
A393R+T2R+S386R,

10 A393R+T490A+V59A+PLASD(N-terminal extension),
S394R+T2Q+A11P,
Y402F+T2R+L66V+S394P,
Y402F+S411V+S119P
Y402F+S411V,
15 Y402F+312Q+S119P+T416H,
S411V+A393R,
E408R+S386N,
E408R+A425T+S465P+T494A,
L410R+A393R,
20 S411V+S119P+402F+A393R,
S411V+S119P+402F+A393R+T416H.

In an embodiment the region mutated is the Region: 445-470.

Preferred sub-regions include one or more of: 445-459, 461-470.

Specific preferred positions contemplated include one or more of:

25 445,446,447,448,449,450,451,452,453,454,455,456,457,458,459,461,
462,463,464,465,466,467,468,469,470.

Specific mutations include one or more of: G447S, G456C/P, S465P.

Preferred combinations of mutations include one or more of:

G447S+S119P,
30 S465P+E408R+A425T+T494A.

Increased specific activity

In a third aspect the invention relates to a variant of a parent glucoamylase with increased specific activity comprising one or more mutation(s) in the following position(s)

35 or region(s) in the amino acid sequence shown in NO: 2:

Region: 1-18,

Region: 40-62,
Region: 93-127,
Region: 170-184,
Region: 200-212,
5 Region: 234-246,
Region: 287-319,
Region: 388-414,

and/or in a corresponding position or region in a homologous glucoamylase which displays at least 60% homology with the amino acid sequences shown in SEQ ID NO: 2,
10 with the exception of the following substitutions: S411G.

In an embodiment the region mutated is the Region: 1-18.

Specific preferred positions contemplated include one or more of:

1,2,3,4,5,6,7,8,9,10,11,12,13,14,15,16,17,18,19.

Specific mutations are: L3N, I18V.

15 Preferred combinations of mutations include one or more of:

I18V+T51S+S56A+V59T+L60A.

In an embodiment the region mutated is the Region: 40-62.

Preferred sub-regions include one or more of: 40-47, 58-62.

Specific preferred positions contemplated include one or more
20 of:40,41,42,43,44,45,46,47,49,51,53,56,58,59,60,61,62.

Specific mutations include one or more of:

S40C,T43R,T51S/D,T53D,S56A/C,V59T,L60A.

Preferred combinations of mutations include one or more of:

T51S+S56A+V59T+L60A+I18V.

25 In an embodiment the region mutated is the Region: 93-127.

In an additional embodiment the sub-region is: Region: 93-124

Preferred sub-regions include one or more of: 93-107, 109-111, 113-115.

Specific preferred positions contemplated include one or more of:

93,94,95,96,97,98,99,100,101,102,103,104,105,106,109,110,111,113,114,115,117,118,1
30 26,127.

Specific mutations include one or more of: N93T, P94V, S95N, D97S, L98S/P, S100T/D, A102S/*, N110T, V111P, D112N, E113M/A, T114S, A115Q/G, Y116F, S119A, G127A.

Preferred combinations of mutations include one or more of:

S119P+Y402F,

35 S119P+Y402F+I189T+Y223F+F227Y.

In an embodiment the region mutated is the Region: 170-184.

Specific mutations include one or more of:

N171R/K/Q/E/W/F/Y,

Q172A/R/N/D/C/E/G/H/I/L/K/F/M/P/S/T/W/Y/V,

T173K/R/S/N/Q,

5 G174A,S,

Y175N/Q/D/E,

D176L

L177I

E180N/M

10 V181I/T

N182R/C/E/G/H/I/L/K/M/P/T/W/Y/V, preferred is E,

G183A

S184D/N/E/Q/L/I/T/R/K.

In an embodiment the region mutated is the Region: 200-212.

15 Preferred sub-regions include: 200-211.

Specific preferred positions contemplated include one or more of:

200,201,202,203,204,205,206,207,208,209,210,211.

Specific mutations include one or more of: A201D, F202L, A203L, T204K, A205R/S, V206L/N, G207N, S208H/T/D, S209T, W212N/A/T.

20 In an embodiment the region mutated is the Region: 234-246.

Preferred sub-regions include one or more of: 234-240, 242-245.

Specific preferred positions contemplated include one or more of:

234,235,236,237,238,239,240,242,243,244,245.

In an embodiment the region mutated is the Region: 287-319.

25 Preferred sub-regions include one or more of: 287-292, 294-301, 313-316.

Specific preferred positions contemplated include one or more of:

287,288,289,290,291,292,294,295,296,297,298,299,300,301,302,

303,307,308,310,311,313,314,315,316,318,319.

Specific mutations include one or more of:

30 S287A/R/N/D/C/Q/E/G/H/I/L/K/M/T/V,

I288L/N/Q,

Y289F,

T290A/R/N/D/C/Q/E/G/H/I/L/K/M/P/S/V,

L291I/D/N,

35 N292D,

D293A/R/N/C/Q/E/G/H/I/L/K/M/S/T/V,

G294A/R/N/D/C/Q/E/H/I/L/K/M/P/S/T/V,
 L295A/R/N/D/C/Q/E/G/H/K/M/S/T/V,
 S296A/R/N/D/C/Q/E/G/H/I/L/K/M/T/V,
 D297A/R/N/C/Q/E/G/H/I/L/K/M/P/S/T/V,
 5 S298A/R/N/D/C/Q/E/G/H/I/L/K/F/M/T/V,
 E299A/R/N/D/C/Q/G/H/I/L/K/M/S/T/V.
 V301T/I,
 A302R/N/D/C/Q/E/G/H/I/L/K/F/M/P/S/T/W/Y/V/, preferred S, V303T/I,
 G304A,
 10 R305A/N/D/C/Q/E/G/H/I/L/F/M/P/S/T/W/Y/V,
 Y306A/R/N/D/C/Q/E/G/H/I/L/K/M/P/S/T/W/V.
 E308A/R/N/D/C/Q/G/H/I/L/K/M/P/S/T/V, preferred Q,
 D309L,
 T310V/S,
 15 Y311N,
 Y312Q/N, preferred is Q,
 N313T/S/G, preferred is S,
 N315Q/E/R.

In an embodiment the region mutated is the Region: 388-414.

20 Preferred sub-regions include one or more of: 397-399, 402-406, 412-414.

Specific preferred positions contemplated include one or more of:

388, 389, 390, 394, 397, 398, 399, 402, 403, 404, 405, 406, 409, 412, 413, 414.

Specific mutations include one or more of: M398L, S399C/Q/T, Y402F, D403S, S405T, E408C/R, S411V, A412C, D414A.

25 Preferred combinations of mutations include one or more of:

Y402F+S119P,

Y402F+S119P+I189T+Y223F+F227Y.

In a preferred embodiment of the invention the regions to be mutated are:

Region: 287-300,

30 Region: 305-319,

and/or corresponding positions or regions in a homologous glucoamylase which displays at least 60% homology with the amino acid sequences shown in SEQ ID NO: 2.

Homology (identity)

35 The homology referred to above of the parent glucoamylase is determined as the degree of identity between two protein sequences indicating a derivation of the first

sequence from the second. The homology may suitably be determined by means of computer programs known in the art such as GAP provided in the GCG program package (Program Manual for the Wisconsin Package, Version 8, August 1994, Genetics Computer Group, 575 Science Drive, Madison, Wisconsin, USA 53711) (Needleman, S.B. and Wunsch, C.D., (1970), Journal of Molecular Biology, 48, p. 443-453). Using Gap with the following settings for polypeptide sequence comparison: Gap creation penalty of 3.0 and Gap extension penalty of 0.1, the mature part of a polypeptide encoded by an analogous DNA sequence of the invention exhibits a degree of identity preferably of at least 60%, such as 70%, at least 80%, at least 90%, more preferably at least 95%, more preferably at least 97%, and most preferably at least 99% with the mature part of the amino acid sequence shown in SEQ ID NO: 2.

Preferably, the parent glucoamylase comprise the amino acid sequences of SEQ ID NO: 2; or allelic variants thereof; or fragments thereof that has glucoamylase activity.

A fragment of SEQ ID NO: 2 is a polypeptide which have one or more amino acids deleted from the amino and/or carboxyl terminus of this amino acid sequence. For instance, the AMG G2 (SEQ ID NO: 2) is a fragment of the *Aspergillus niger* G1 glucoamylase (Boel et al. (1984), EMBO J. 3 (5), p. 1097-1102) having glucoamylase activity. An allelic variant denotes any of two or more alternative forms of a gene occupying the same chromosomal locus. Allelic variation arises naturally through mutation, and may result in polymorphism within populations. Gene mutations can be silent (no change in the encoded polypeptide) or may encode polypeptides having altered amino acid sequences. An allelic variant of a polypeptide is a polypeptide encoded by an allelic variant of a gene.

The amino acid sequences of homologous parent glucoamylases may differ from the amino acid sequence of SEQ ID NO: 2 by an insertion or deletion of one or more amino acid residues and/or the substitution of one or more amino acid residues by different amino acid residues. Preferably, amino acid changes are of a minor nature, that is conservative amino acid substitutions that do not significantly affect the folding and/or activity of the protein; small deletions, typically of one to about 30 amino acids; small amino- or carboxyl-terminal extensions, such as an amino-terminal methionine residue; a small linker peptide of up to about 20-25 residues; or a small extension that facilitates purification by changing net charge or another function, such as a poly-histidine tract, an antigenic epitope or a binding domain.

In another embodiment, the isolated parent glucoamylase is encoded by a nucleic acid sequence which hybridises under very low stringency conditions, preferably low stringency conditions, more preferably medium stringency conditions, more preferably

medium-high stringency conditions, even more preferably high stringency conditions, and most preferably very high stringency conditions with a nucleic acid probe which hybridises under the same conditions with (i) the nucleic acid sequence of SEQ ID NO: 1, (ii) the cDNA sequence of SEQ ID NO:1, (iii) a sub-sequence of (i) or (ii), or (iv) a complementary strand of (i), (ii), or (iii) (J. Sambrook, E.F. Fritsch, and T. Maniatus, 5 1989, *Molecular Cloning, A Laboratory Manual*, 2d edition, Cold Spring Harbor, New York). The sub-sequence of SEQ ID NO: 1 may be at least 100 nucleotides or preferably at least 200 nucleotides. Moreover, the sub-sequence may encode a polypeptide fragment which has glucoamylase activity. The parent polypeptides may also be allelic variants or fragments of the polypeptides that have glucoamylase activity. 10

The nucleic acid sequence of SEQ ID NO: 1 or a subsequence thereof, as well as the amino acid sequence of SEQ ID NO: 2, or a fragment thereof, may be used to design a nucleic acid probe to identify and clone DNA encoding polypeptides having glucoamylase activity, from strains of different genera or species according to methods well known in the art. In particular, such probes can be used for hybridization with the genomic or cDNA of the genus or species of interest, following standard Southern blotting procedures, in order to identify and isolate the corresponding gene therein. Such probes can be considerably shorter than the entire sequence, but should be at least 15, preferably at least 25, and more preferably at least 35 nucleotides in length. 15 Longer probes can also be used. Both DNA and RNA probes can be used. The probes are typically labeled for detecting the corresponding gene (for example, with ³²P, ³H, ³⁵S, biotin, or avidin). Such probes are encompassed by the present invention. 20

Thus, a genomic DNA or cDNA library prepared from such other organisms may be screened for DNA which hybridizes with the probes described above and which encodes a polypeptide having glucoamylase. Genomic or other DNA from such other organisms may be separated by agarose or polyacrylamide gel electrophoresis, or other separation techniques. DNA from the libraries or the separated DNA may be transferred to and immobilised on nitrocellulose or other suitable carrier material. In order to identify a clone or DNA which is homologous with SEQ ID NO: 1, or sub-sequences thereof, the carrier material is used in a Southern blot. For purposes of the present invention, hybridisation indicates that the nucleic acid sequence hybridises to a nucleic acid probe corresponding to the nucleic acid sequence shown in SEQ ID NO: 1 its complementary strand, or a sub-sequence thereof, under very low to very high stringency conditions. Molecules to which the nucleic acid probe hybridises under 35 these conditions are detected using X-ray film.

For long probes of at least 100 nucleotides in length, the carrier material is finally washed three times each for 15 minutes using 2 x SSC, 0.2% SDS preferably at least at 45°C (very low stringency), more preferably at least at 50°C (low stringency), more preferably at least at 55°C (medium stringency), more preferably at least at 60°C (medium-high stringency), even more preferably at least at 65°C (high stringency), and most preferably at least at 70°C (very high stringency).

For short probes which are about 15 nucleotides to about 70 nucleotides in length, stringency conditions are defined as prehybridization, hybridisation, and washing post-hybridization at 5°C to 10°C below the calculated T_m using the calculation according to Bolton and McCarthy (1962, *Proceedings of the National Academy of Sciences USA* 48:1390) in 0.9 M NaCl, 0.09 M Tris-HCl pH 7.6, 6 mM EDTA, 0.5% NP-40, 1X Denhardt's solution, 1 mM sodium pyrophosphate, 1 mM sodium monobasic phosphate, 0.1 mM ATP, and 0.2 mg of yeast RNA per ml following standard Southern blotting procedures.

For short probes which are about 15 nucleotides to about 70 nucleotides in length, the carrier material is washed once in 6X SSC plus 0.1% SDS for 15 minutes and twice each for 15 minutes using 6X SSC at 5°C to 10°C below the calculated T_m .

The present invention also relates to isolated nucleic acid sequences produced by (a) hybridising a DNA under very low, low, medium, medium-high, high, or very high stringency conditions with the sequence of SEQ ID NO:1, or its complementary strand, or a sub-sequence thereof; and (b) isolating the nucleic acid sequence. The sub-sequence is preferably a sequence of at least 100 nucleotides such as a sequence which encodes a polypeptide fragment which has glucoamylase activity.

Contemplated parent glucoamylases have at least 20%, preferably at least 40%, more preferably at least 60%, even more preferably at least 80%, even more preferably at least 90%, and most preferably at least 100% of the glucoamylase activity of the mature polypeptide of SEQ ID NO: 2.

In a preferred embodiment the variant of the invention has improved thermal stability and/or increased specific activity, preferably within the temperature interval from about 60-80°C, preferably 63-75°C, preferably at a pH of 4-5, in particular 4.2-4.7, using maltodextrin as the substrate.

In another preferred embodiment a variant of the invention is used for, e.g., alcohol fermentation.

In a preferred embodiment the parent glucoamylase is the *Aspergillus niger* G1 glucoamylase (Boel et al. (1984), EMBO J. 3 (5), p. 1097-1102. The parent glucoamylase may be a truncated glucoamylase, e.g., the AMG G2 glucoamylase.

Cloning a DNA Sequence Encoding A Parent Glucoamylase

The DNA sequence encoding a parent glucoamylase may be isolated from any cell or microorganism producing the glucoamylase in question, using various methods well known in the art. First, a genomic DNA and/or cDNA library should be constructed using chromosomal DNA or messenger RNA from the organism that produces the glucoamylase to be studied. Then, if the amino acid sequence of the glucoamylase is known, labeled oligonucleotide probes may be synthesized and used to identify glucoamylase-encoding clones from a genomic library prepared from the organism in question. Alternatively, a labelled oligonucleotide probe containing sequences homologous to another known glucoamylase gene could be used as a probe to identify glucoamylase-encoding clones, using hybridization and washing conditions of very low to very high stringency. This is described above.

Yet another method for identifying glucoamylase-encoding clones would involve inserting fragments of genomic DNA into an expression vector, such as a plasmid, transforming glucoamylase-negative bacteria with the resulting genomic DNA library, and then plating the transformed bacteria onto agar containing a substrate for glucoamylase (*i.e.*, maltose), thereby allowing clones expressing the glucoamylase to be identified.

Alternatively, the DNA sequence encoding the enzyme may be prepared synthetically by established standard methods, *e.g.* the phosphoroamidite method described S.L. Beaucage and M.H. Caruthers, (1981), Tetrahedron Letters 22, p. 1859-1869, or the method described by Matthes et al., (1984), EMBO J. 3, p. 801-805. In the phosphoroamidite method, oligonucleotides are synthesized, *e.g.*, in an automatic DNA synthesizer, purified, annealed, ligated and cloned in appropriate vectors.

Finally, the DNA sequence may be of mixed genomic and synthetic origin, mixed synthetic and cDNA origin or mixed genomic and cDNA origin, prepared by ligating fragments of synthetic, genomic or cDNA origin (as appropriate, the fragments corresponding to various parts of the entire DNA sequence), in accordance with standard techniques. The DNA sequence may also be prepared by polymerase chain reaction (PCR) using specific primers, for instance as described in US 4,683,202 or R.K. Saiki et al., (1988), Science 239, 1988, pp. 487-491.

Site-directed mutagenesis

Once a glucoamylase-encoding DNA sequence has been isolated, and desirable sites for mutation identified, mutations may be introduced using synthetic oligonucleotides. These oligonucleotides contain nucleotide sequences flanking the

desired mutation sites. In a specific method, a single-stranded gap of DNA, the glucoamylase-encoding sequence, is created in a vector carrying the glucoamylase gene. Then the synthetic nucleotide, bearing the desired mutation, is annealed to a homologous portion of the single-stranded DNA. The remaining gap is then filled in with DNA polymerase I (Klenow fragment) and the construct is ligated using T4 ligase. A specific example of this method is described in Morinaga et al., (1984), Biotechnology 2, p. 646-639. US 4,760,025 discloses the introduction of oligonucleotides encoding multiple mutations by performing minor alterations of the cassette. However, an even greater variety of mutations can be introduced at any one time by the Morinaga method, because a multitude of oligonucleotides, of various lengths, can be introduced.

Another method for introducing mutations into glucoamylase-encoding DNA sequences is described in Nelson and Long, (1989), Analytical Biochemistry 180, p. 147-151. It involves the 3-step generation of a PCR fragment containing the desired mutation introduced by using a chemically synthesized DNA strand as one of the primers in the PCR reactions. From the PCR-generated fragment, a DNA fragment carrying the mutation may be isolated by cleavage with restriction endonucleases and reinserted into an expression plasmid.

Further, Sierks. et al., (1989), Protein Eng., 2, 621-625; Sierks et al., (1990), Protein Eng. vol. 3, 193-198; also describes site-directed mutagenesis in an *Aspergillus* glucoamylase.

Random Mutagenesis

Random mutagenesis is suitably performed either as localized or region-specific random mutagenesis in at least three parts of the gene translating to the amino acid sequence shown in question, or within the whole gene.

The random mutagenesis of a DNA sequence encoding a parent glucoamylase may be conveniently performed by use of any method known in the art.

In relation to the above, a further aspect of the present invention relates to a method for generating a variant of a parent glucoamylase, wherein the variant exhibits increased thermal stability relative to the parent, the method comprising:

- (a) subjecting a DNA sequence encoding the parent glucoamylase to random mutagenesis,
- (b) expressing the mutated DNA sequence obtained in step (a) in a host cell, and
- (c) screening for host cells expressing a glucoamylase variant which has an altered property (i.e. thermal stability) relative to the parent glucoamylase.

Step (a) of the above method of the invention is preferably performed using doped primers, as described in the working examples herein (*vide infra*).

For instance, the random mutagenesis may be performed by use of a suitable physical or chemical mutagenizing agent, by use of a suitable oligonucleotide, or by
5 subjecting the DNA sequence to PCR generated mutagenesis. Furthermore, the random mutagenesis may be performed by use of any combination of these mutagenizing agents. The mutagenizing agent may, *e.g.*, be one which induces transitions, transversions, inversions, scrambling, deletions, and/or insertions.

Examples of a physical or chemical mutagenizing agent suitable for the present
10 purpose include ultraviolet (UV) ir-radiation, hydroxylamine, N-methyl-N'-nitro-N-nitrosoguanidine (MNNG), O-methyl hydroxylamine, nitrous acid, ethyl methane sulphonate (EMS), sodium bisulphite, formic acid, and nucleotide analogues. When such agents are used, the mutagenesis is typically performed by incubating the DNA sequence encoding the parent enzyme to be mutagenized in the presence of the mutagenizing
15 agent of choice under suitable conditions for the mutagenesis to take place, and selecting for mutated DNA having the desired properties.

When the mutagenesis is performed by the use of an oligonucleotide, the oligonucleotide may be doped or spiked with the three non-parent nucleotides during the synthesis of the oligonucleotide at the positions which are to be changed. The doping or
20 spiking may be done so that codons for unwanted amino acids are avoided. The doped or spiked oligonucleotide can be incorporated into the DNA encoding the glucoamylase enzyme by any published technique, using, *e.g.*, PCR, LCR or any DNA polymerase and ligase as deemed appropriate.

Preferably, the doping is carried out using "constant random doping", in which the
25 percentage of wild-type and mutation in each position is predefined. Furthermore, the doping may be directed toward a preference for the introduction of certain nucleotides, and thereby a preference for the introduction of one or more specific amino acid residues. The doping may be made, *e.g.*, so as to allow for the introduction of 90% wild type and 10% mutations in each position. An additional consideration in the choice of a doping
30 scheme is based on genetic as well as protein-structural constraints. The doping scheme may be made by using the DOPE program which, *inter alia*, ensures that introduction of stop codons is avoided.

When PCR-generated mutagenesis is used, either a chemically treated or non-treated gene encoding a parent glucoamylase is subjected to PCR under conditions that
35 increase the mis-incorporation of nucleotides (Deshler 1992; Leung et al., Technique, Vol.1, 1989, pp. 11-15).

A mutator strain of *E. coli* (Fowler et al., Molec. Gen. Genet., 133, 1974, pp. 179-191), *S. cerevisiae* or any other microbial organism may be used for the random mutagenesis of the DNA encoding the glucoamylase by, e.g., transforming a plasmid containing the parent glycosylase into the mutator strain, growing the mutator strain with the plasmid and isolating the mutated plasmid from the mutator strain. The mutated plasmid may be subsequently transformed into the expression organism.

The DNA sequence to be mutagenized may be conveniently present in a genomic or cDNA library prepared from an organism expressing the parent glucoamylase. Alternatively, the DNA sequence may be present on a suitable vector such as a plasmid or a bacteriophage, which as such may be incubated with or otherwise exposed to the mutagenising agent. The DNA to be mutagenized may also be present in a host cell either by being integrated in the genome of said cell or by being present on a vector harboured in the cell. Finally, the DNA to be mutagenized may be in isolated form. It will be understood that the DNA sequence to be subjected to random mutagenesis is preferably a cDNA or a genomic DNA sequence.

In some cases it may be convenient to amplify the mutated DNA sequence prior to performing the expression step b) or the screening step c). Such amplification may be performed in accordance with methods known in the art, the presently preferred method being PCR-generated amplification using oligonucleotide primers prepared on the basis of the DNA or amino acid sequence of the parent enzyme.

Subsequent to the incubation with or exposure to the mutagenising agent, the mutated DNA is expressed by culturing a suitable host cell carrying the DNA sequence under conditions allowing expression to take place. The host cell used for this purpose may be one which has been transformed with the mutated DNA sequence, optionally present on a vector, or one which carried the DNA sequence encoding the parent enzyme during the mutagenesis treatment. Examples of suitable host cells are the following: gram positive bacteria such as *Bacillus subtilis*, *Bacillus licheniformis*, *Bacillus lentus*, *Bacillus brevis*, *Bacillus stearothermophilus*, *Bacillus alkalophilus*, *Bacillus amyloliquefaciens*, *Bacillus coagulans*, *Bacillus circulans*, *Bacillus lautus*, *Bacillus megaterium*, *Bacillus thuringiensis*, *Streptomyces lividans* or *Streptomyces murinus*; and gram-negative bacteria such as *E. coli*.

The mutated DNA sequence may further comprise a DNA sequence encoding functions permitting expression of the mutated DNA sequence.

35 Localized random mutagenesis

The random mutagenesis may be advantageously localized to a part of the parent glucoamylase in question. This may, e.g., be advantageous when certain regions of the enzyme have been identified to be of particular importance for a given property of the enzyme, and when modified are expected to result in a variant having improved properties. Such regions may normally be identified when the tertiary structure of the parent enzyme has been elucidated and related to the function of the enzyme.

The localized, or region-specific, random mutagenesis is conveniently performed by use of PCR generated mutagenesis techniques as described above or any other suitable technique known in the art. Alternatively, the DNA sequence encoding the part of the DNA sequence to be modified may be isolated, e.g., by insertion into a suitable vector, and said part may be subsequently subjected to mutagenesis by use of any of the mutagenesis methods discussed above.

Alternative methods for providing variants of the invention include gene shuffling e.g. as described in WO 95/22625 (from Affymax Technologies N.V.) or in WO 96/00343 (from Novo Nordisk A/S).

Expression Of Glucoamylase Variants

According to the invention, a DNA sequence encoding the variant produced by methods described above, or by any alternative methods known in the art, can be expressed, in enzyme form, using an expression vector which typically includes control sequences encoding a promoter, operator, ribosome binding site, translation initiation signal, and, optionally, a repressor gene or various activator genes.

Expression vector

The recombinant expression vector carrying the DNA sequence encoding a glucoamylase variant of the invention may be any vector which may conveniently be subjected to recombinant DNA procedures, and the choice of vector will often depend on the host cell into which it is to be introduced. The vector may be one which, when introduced into a host cell, is integrated into the host cell genome and replicated together with the chromosome(s) into which it has been integrated. Examples of suitable expression vectors include pMT838.

Promoter

In the vector, the DNA sequence should be operably connected to a suitable promoter sequence. The promoter may be any DNA sequence which shows transcrip-

tional activity in the host cell of choice and may be derived from genes encoding proteins either homologous or heterologous to the host cell.

Examples of suitable promoters for directing the transcription of the DNA sequence encoding a glucoamylase variant of the invention, especially in a bacterial host, are the promoter of the *lac* operon of *E.coli*, the *Streptomyces coelicolor* agarase gene *dagA* promoters, the promoters of the *Bacillus licheniformis* α -amylase gene (*amyL*), the promoters of the *Bacillus stearothermophilus* maltogenic amylase gene (*amyM*), the promoters of the *Bacillus amyloliquefaciens* α -amylase (*amyQ*), the promoters of the *Bacillus subtilis* *xylA* and *xylB* genes etc. For transcription in a fungal host, examples of useful promoters are those derived from the gene encoding *A. oryzae* TAKA amylase, the TPI (triose phosphate isomerase) promoter from *S. cerevisiae* (Alber et al. (1982), J. Mol. Appl. Genet 1, p. 419-434, *Rhizomucor miehei* aspartic proteinase, *A. niger* neutral α -amylase, *A. niger* acid stable α -amylase, *A. niger* glucoamylase, *Rhizomucor miehei* lipase, *A. oryzae* alkaline protease, *A. oryzae* triose phosphate isomerase or *A. nidulans* acetamidase.

Expression vector

The expression vector of the invention may also comprise a suitable transcription terminator and, in eukaryotes, polyadenylation sequences operably connected to the DNA sequence encoding the α -amylase variant of the invention. Termination and polyadenylation sequences may suitably be derived from the same sources as the promoter.

The vector may further comprise a DNA sequence enabling the vector to replicate in the host cell in question. Examples of such sequences are the origins of replication of plasmids pUC19, pACYC177, pUB110, pE194, pAMB1 and pIJ702.

The vector may also comprise a selectable marker, e.g. a gene the product of which complements a defect in the host cell, such as the *dal* genes from *B. subtilis* or *B. licheniformis*, or one which confers antibiotic resistance such as ampicillin, kanamycin, chloramphenicol or tetracyclin resistance. Furthermore, the vector may comprise *Aspergillus* selection markers such as *amdS*, *argB*, *niaD* and *sC*, a marker giving rise to hygromycin resistance, or the selection may be accomplished by co-transformation, e.g. as described in WO 91/17243.

The procedures used to ligate the DNA construct of the invention encoding a glucoamylase variant, the promoter, terminator and other elements, respectively, and to insert them into suitable vectors containing the information necessary for replication, are well known to persons skilled in the art (cf., for instance, Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd Ed., Cold Spring Harbor, 1989).

Host Cells

The cell of the invention, either comprising a DNA construct or an expression vector of the invention as defined above, is advantageously used as a host cell in the recombinant production of a glucoamylase variant of the invention. The cell may be transformed with the DNA construct of the invention encoding the variant, conveniently by integrating the DNA construct (in one or more copies) in the host chromosome. This integration is generally considered to be an advantage as the DNA sequence is more likely to be stably maintained in the cell. Integration of the DNA constructs into the host chromosome may be performed according to conventional methods, e.g. by homologous or heterologous recombination. Alternatively, the cell may be transformed with an expression vector as described above in connection with the different types of host cells.

The cell of the invention may be a cell of a higher organism such as a mammal or an insect, but is preferably a microbial cell, e.g., a bacterial or a fungal (including yeast) cell.

Examples of suitable bacteria are Gram positive bacteria such as *Bacillus subtilis*, *Bacillus licheniformis*, *Bacillus lentus*, *Bacillus brevis*, *Bacillus stearothermophilus*, *Bacillus alkalophilus*, *Bacillus amyloliquefaciens*, *Bacillus coagulans*, *Bacillus circulans*, *Bacillus lautus*, *Bacillus megaterium*, *Bacillus thuringiensis*, or *Streptomyces lividans* or *Streptomyces murinus*, or gramnegative bacteria such as *E.coli*. The transformation of the bacteria may, for instance, be effected by protoplast transformation or by using competent cells in a manner known *per se*.

The yeast organism may favorably be selected from a species of *Saccharomyces* or *Schizosaccharomyces*, e.g. *Saccharomyces cerevisiae*.

The host cell may also be a filamentous fungus e.g. a strain belonging to a species of *Aspergillus*, most preferably *Aspergillus oryzae* or *Aspergillus niger*, or a strain of *Fusarium*, such as a strain of *Fusarium oxysporium*, *Fusarium graminearum* (in the perfect state named *Gibberella zeae*, previously *Sphaeria zeae*, synonym with *Gibberella roseum* and *Gibberella roseum* f. sp. *cerealis*), or *Fusarium sulphureum* (in the perfect state named *Gibberella puricaris*, synonym with *Fusarium trichothecioides*, *Fusarium bactridioides*, *Fusarium sambucium*, *Fusarium roseum*, and *Fusarium roseum* var. *graminearum*), *Fusarium cerealis* (synonym with *Fusarium crockwellense*), or *Fusarium venenatum*.

In a preferred embodiment of the invention the host cell is a protease deficient or protease minus strain.

This may for instance be the protease deficient strain *Aspergillus oryzae* JaL 125 having the alkaline protease gene named "alp" deleted. This strain is described in WO 97/35956 (Novo Nordisk).

Filamentous fungi cells may be transformed by a process involving protoplast formation and transformation of the protoplasts followed by regeneration of the cell wall in a manner known per se. The use of *Aspergillus* as a host micro-organism is described in EP 238 023 (Novo Nordisk A/S), the contents of which are hereby incorporated by reference.

Method Of Producing A Glucoamylase Variant

In a yet further aspect, the present invention relates to a method of producing a glucoamylase variant of the invention, which method comprises cultivating a host cell under conditions conducive to the production of the variant and recovering the variant from the cells and/or culture medium.

The medium used to cultivate the cells may be any conventional medium suitable for growing the host cell in question and obtaining expression of the glucoamylase variant of the invention. Suitable media are available from commercial suppliers or may be prepared according to published recipes (e.g. as described in catalogues of the American Type Culture Collection).

The glucoamylase variant secreted from the host cells may conveniently be recovered from the culture medium by well-known procedures, including separating the cells from the medium by centrifugation or filtration, and precipitating proteinaceous components of the medium by means of a salt such as ammonium sulphate, followed by the use of chromatographic procedures such as ion exchange chromatography, affinity chromatography, or the like.

Starch Conversion

The present invention provides a method of using glucoamylase variants of the invention for producing glucose and the like from starch. Generally, the method includes the steps of partially hydrolyzing precursor starch in the presence of α -amylase and then further hydrolyzing the release of D-glucose from the non-reducing ends of the starch or related oligo- and polysaccharide molecules in the presence of glucoamylase by cleaving α -(1 \rightarrow 4) and α -(1 \rightarrow 6) glucosidic bonds.

The partial hydrolysis of the precursor starch utilizing α -amylase provides an initial breakdown of the starch molecules by hydrolyzing internal α -(1 \rightarrow 4)-linkages. In commercial applications, the initial hydrolysis using α -amylase is run at a temperature of

approximately 105°C. A very high starch concentration is processed, usually 30% to 40% solids. The initial hydrolysis is usually carried out for five minutes at this elevated temperature. The partially hydrolyzed starch can then be transferred to a second tank and incubated for approximately 1-2 hour at a temperature of 85° to 98°C to derive a dextrose
5 equivalent (D.E.) of 10 to 15.

The step of further hydrolyzing the release of D-glucose from the non-reducing ends of the starch or related oligo- and polysaccharides molecules in the presence of glucoamylase is normally carried out in a separate tank at a reduced temperature between 30° and 62°C. Preferably the temperature of the substrate liquid is dropped to
10 between 55° and 60°C. The pH of the solution is dropped from about 5.5 to 6.5 to a range between 3 and 5.5. Preferably, the pH of the solution is 4 to 4.5. The glucoamylase is added to the solution and the reaction is carried out for 24-72 hours, preferably 36-48 hours.

By using a thermostable glucoamylase variant of the invention saccharification
15 processes may be carried out at a higher temperature than traditional batch saccharification processes. According to the invention saccharification may be carried out at temperatures in the range from above 60-80°C, preferably 63-75°C. This apply both for traditional batch processes (described above) and for continuous saccharification processes.

20 Actually, continuous saccharification processes including one or more membrane separation steps, *i.e.* filtration steps, must be carried out at temperatures of above 60°C to be able to maintain a reasonably high flux over the membrane or to minimize microbial contamination. Therefore, the thermostable variants of the invention provides the possibility of carrying out large scale continuous saccharification processes at a fair price
25 and/or at a lower enzyme protein dosage within a period of time acceptable for industrial saccharification processes. According to the invention the saccharification time may even be shortened.

The activity of the glucoamylase variant (*e.g.* AMG variant) of the invention is generally substantially higher at temperatures between 60°C-80°C than at the traditionally
30 used temperature between 30-60°C. Therefore, by increasing the temperature at which the glucoamylase operates the saccharification process may be carried out within a shorter period of time.

Further, by improving the thermal stability the $T_{1/2}$ (half-time, as defined in the "Materials and Methods" section) is improved. As the thermal stability of the glucoamylase
35 variants of the invention is improved a minor amount of glucoamylase need to be added to replace the glucoamylase being inactivated during the saccharification process. More

glucoamylase is maintained active during saccharification process according to the present invention. Furthermore, the risk of microbial contamination is also reduced when carrying the saccharification process at temperature above 63°C.

The glucose yield from a typical saccharification trial with glucoamylase, acid
5 amylase and pullulanase is 95.5-96.5%. The remaining carbohydrates typically consists of 1% maltose, 1.5-2% isomaltose and 1-1.5% higher oligosaccharides. The disaccharides are produced since the glucoamylase at high concentrations of glucose and high dry-solid levels has a tendency to form reversion products.

A glucoamylase with an increased specific activity towards saccharides present in
10 the solution after liquefaction and saccharides formed during saccharification would be an advantage as a reduced enzyme protein dosage or a shorter process time then could be used. In general, the glucoamylase has a preference for substrates consisting of longer saccharides compared to short chain saccharides and the specific activity towards e.g. maltoheptaose is therefore approximately 6 times higher than towards maltose. An
15 increased specific activity towards short chain saccharides such as maltose (without reducing the activity towards oligosaccharides) would therefore also permit using a lower enzyme dosage and/or shorter process time.

Furthermore, a higher glucose yield can be obtained with a glucoamylase variant with an increased alpha-1,4 hydrolytic activity (if the alpha-1,6 activity is unchanged or
20 even decreased), since a reduced amount of enzyme protein is being used, and alpha-1,6 reversion product formation therefore is decreased (less isomaltose).

The specific activity may be measured using the method described in the "Materials & Methods" section at 37°C or 60°C.

Example of saccharification process wherein the glucoamylase variants of the
25 invention may be used include the processes described in JP 3-224493; JP 1-191693; JP 62-272987; EP 452,238, and WO 99/27124 (all references are hereby incorporated by reference).

In a further aspect the invention relates to a method of saccharifying a liquefied starch solution, comprising the steps

- 30 (i) a saccharification step during which step one or more enzymatic saccharification stages takes place, and the subsequent step of
(ii) one or more high temperature membrane separation steps
wherein the enzymatic saccharification is carried out using a thermostable glucoamylase variant of the invention.

35 The glucoamylase variant(s) of the invention may be used in the present inventive process in combination with an enzyme that hydrolyzes only α -(1 \rightarrow 6)-glucosidic bonds in

molecules with at least four glucosyl residues. Preferentially, the glucoamylase variant of the invention can be used in combination with pullulanase or isoamylase. The use of isoamylase and pullulanase for debranching, the molecular properties of the enzymes, and the potential use of the enzymes with glucoamylase is set forth in G.M.A. van Beynum et al., Starch Conversion Technology, Marcel Dekker, New York, 1985, 101-142.

In a further aspect the invention relates to the use of a glucoamylase variant of the invention in a starch conversion process.

Further, the glucoamylase variant of the invention may be used in a continuous starch conversion process including a continuous saccharification step.

The glucoamylase variants of the invention may also be used in immobilised form. This is suitable and often used for producing speciality syrups, such as maltose syrups, and further for the raffinate stream of oligosaccharides in connection with the production of fructose syrups.

The glucoamylase of the invention may also be used in a process for producing ethanol for fuel or beverage or may be used in a fermentation process for producing organic compounds, such as citric acid, ascorbic acid, lysine, glutamic acid.

MATERIALS & METHODS

MATERIALS:

Enzymes:

AMG G1: *Aspergillus niger* glucoamylase G1 disclosed in Boel et al. (1984), EMBO J. 3 (5), 1097-1102, and SEQ ID NO: 13, available from Novo Nordisk.

AMG G2: Truncated *Aspergillus niger* glucoamylase G1 shown in SEQ ID NO: 2, available from Novo Nordisk)

Solutions:

Buffer: 0.05M sodium acetate (6.8g in 1 l milli-Q-water), pH 4.5

Stop solution: 0.4M NaOH

GOD-perid, 124036, Boehringer Mannheim

Substrate:

Maltose: 29mM (1g maltose in 100ml 50mM sodium acetate, pH 4.5) (Sigma)

Maltoheptaose: 10 mM, 115 mg/10 ml (Sigma)

Host cell:

A. oryzae JaL 125: *Aspergillus oryzae* IFO 4177 available from Institute for Fermentation, Osaka; 17-25 Juso Hammachi 2-Chome Yodogawa-ku, Osaka, Japan, having the alkaline protease gene named "alp" (described by Murakami K et al., (1991), Agric. Biol. Chem. 55, p. 2807-2811) deleted by a one step gene replacement method
5 (described by G. May in "Applied Molecular Genetics of Filamentous Fungi" (1992), p. 1-25. Eds. J. R. Kinghorn and G. Turner; Blackie Academic and Professional), using the *A. oryzae* pyrG gene as marker. Strain JaL 125 is further disclosed in WO 97/35956 (Novo Nordisk).

10 **Micro-organisms:**

Strain: *Saccharomyces cerevisiae* YNG318: MAT α leu2- Δ 2 ura3-52 his4-539 pep4- Δ 1[cir+]

Plasmids:

15 pCAMG91: see Figure 1. Plasmid comprising the *Aspergillus niger* G1 glucoamylase (AMG G1). The construction of pCAMG91 is described in Boel et al. (1984), EMBO J. 3 (7) p.1581-1585.

pMT838: Plasmid encoding the truncated *Aspergillus niger* glucoamylase G2 (SEQ ID NO: 2).
20

pJSO026 (*S. cerevisiae* expression plasmid)(J.S.Okkels, (1996)"A URA3-promoter deletion in a pYES vector increases the expression level of a fungal lipase in *Saccharomyces cerevisiae*. Recombinant DNA Biotechnology III: The Integration of
25 Biological and Engineering Sciences, vol. 782 of the Annals of the New York Academy of Sciences) More specifically, the expression plasmid pJSO37, is derived from pYES 2.0 by replacing the inducible GAL1-promoter of pYES 2.0 with the constitutively expressed TPI (triose phosphate isomerase)-promoter from *Saccharomyces cerevisiae* (Albert and Karwasaki, (1982), J. Mol. Appl Genet., 1, 419-434), and deleting a part of the URA3
30 promoter.

METHODS:

Transformation of *Saccharomyces cerevisiae* YNG318

The DNA fragments and the opened vectors are mixed and transformed into the
35 yeast *Saccharomyces cerevisiae* YNG318 by standard methods.

Determining Specific Activity As k_{cat} (sec.⁻¹).

750 microL substrate (1% maltose, 50 mM Sodium acetat, pH 4.3) is incubated 5 minutes at selected temperature, such as 37°C or 60°C.

50 microL enzyme diluted in sodium acetate is added.

Aliquots of 100 microL are removed after 0, 3, 6, 9 and 12 minutes and transferred to

5 100 microL 0.4 M Sodium hydroxide to stop the reaction. A blank is included.

20 microL is transferred to a Micro titre plates and 200 microL GOD-Perid solution is added. Absorbance is measured at 650 nm after 30 minutes incubation at room temperature.

Glucose is used as standard and the specific activity is calculated as k_{cat} (sec.⁻¹).

10

Determination Of AGU Activity and As AGU/mg

One Novo Amyloglucosidase Unit (AGU) is defined as the amount of enzyme which hydrolyzes 1 micromole maltose per minute at 37°C and pH 4.3. A detailed description of the analytical method (AEL-SM-0131) is available on request from Novo Nordisk .

15 The activity is determined as AGU/ml by a method modified after (AEL-SM-0131) using the Glucose GOD-Perid kit from Boehringer Mannheim, 124036. Standard: AMG-standard, batch 7-1195, 195 AGU/ml.

375 microL substrate (1% maltose in 50 mM Sodium acetate, pH 4.3) is incubated 5 minutes at 37°C. 25 microL enzyme diluted in sodium acetate is added. The reaction is stopped after 10 minutes by adding 100 microL 0.25 M NaOH. 20 microL is transferred to a 96 well microtitre plate and 200 microL GOD-Perid solution is added. After 30 minutes at room temperature, the absorbance is measured at 650 nm and the activity calculated in AGU/ml from the AMG-standard.

20 The specific activity in AGU/mg is then calculated from the activity (AGU/ml) 25 divided with the protein concentration (mg/ml).

Transformation Of *Aspergillus oryzae* (general procedure)

100 ml of YPD (Sherman et al., (1981), Methods in Yeast Genetics, Cold Spring Harbor Laboratory) are inoculated with spores of *A. oryzae* and incubated with shaking 30 for about 24 hours. The mycelium is harvested by filtration through miracloth and washed with 200 ml of 0.6 M MgSO₄. The mycelium is suspended in 15 ml of 1.2 M MgSO₄, 10 mM NaH₂PO₄, pH 5.8. The suspension is cooled on ice and 1 ml of buffer containing 120 mg of NovozymTM 234 is added. After 5 min., 1 ml of 12 mg/ml BSA (Sigma type H25) is added and incubation with gentle agitation continued for 1.5-2.5 35 hours at 37C until a large number of protoplasts is visible in a sample inspected under the microscope.

The suspension is filtered through miracloth, the filtrate transferred to a sterile tube and overlaid with 5 ml of 0.6 M sorbitol, 100 mM Tris-HCl, pH 7.0. Centrifugation is performed for 15 min. at 1000 g and the protoplasts are collected from the top of the MgSO₄ cushion. 2 volumes of STC (1.2 M sorbitol, 10 mM Tris-HCl, pH 7.5, 10 mM CaCl₂) are added to the protoplast suspension and the mixture is centrifugated for 5 min. at 1000 g. The protoplast pellet is resuspended in 3 ml of STC and repelleted. This is repeated. Finally, the protoplasts are resuspended in 0.2-1 ml of STC.

100 µl of protoplast suspension are mixed with 5-25 µg of p3SR2 (an *A. nidulans* amdS gene carrying plasmid described in Hynes et al., Mol. and Cel. Biol., Vol. 3, No. 8, 1430-1439, Aug. 1983) in 10 µl of STC. The mixture is left at room temperature for 25 min. 0.2 ml of 60% PEG 4000 (BDH 29576), 10 mM CaCl₂ and 10 mM Tris-HCl, pH 7.5 is added and carefully mixed (twice) and finally 0.85 ml of the same solution are added and carefully mixed. The mixture is left at room temperature for 25 min., spun at 2.500 g for 15 min. and the pellet is resuspended in 2 ml of 1.2M sorbitol. After one more sedimentation the protoplasts are spread on minimal plates (Cove, (1966), Biochem. Biophys. Acta 113, 51-56) containing 1.0 M sucrose, pH 7.0, 10 mM acetamide as nitrogen source and 20 mM CsCl to inhibit background growth. After incubation for 4-7 days at 37°C spores are picked, suspended in sterile water and spread for single colonies. This procedure is repeated and spores of a single colony after the second re-isolation are stored as a defined transformant.

Fed Batch Fermentation

Fed batch fermentation is performed in a medium comprising maltodextrin as a carbon source, urea as a nitrogen source and yeast extract. The fed batch fermentation is performed by inoculating a shake flask culture of *A. oryzae* host cells in question into a medium comprising 3.5% of the carbon source and 0.5% of the nitrogen source. After 24 hours of cultivation at pH 5.0 and 34°C the continuous supply of additional carbon and nitrogen sources are initiated. The carbon source is kept as the limiting factor and it is secured that oxygen is present in excess. The fed batch cultivation is continued for 4 days, after which the enzymes can be recovered by centrifugation, ultrafiltration, clear filtration and germ filtration.

Purification

The culture broth is filtrated and added ammoniumsulphate (AMS) to a concentration of 1.7 M AMS and pH is adjusted to pH 5. Precipitated material is removed by centrifugation and the solution containing glucoamylase activity is applied

on a Toyo Pearl Butyl column previously equilibrated in 1.7 M AMS, 20 mM sodium acetate, pH 5. Unbound material is washed out with the equilibration buffer. Bound proteins are eluted with 10 mM sodium acetate, pH 4.5 using a linear gradient from 1.7 - 0 M AMS over 10 column volumes. Glucoamylase containing fractions are collected and dialysed against 20 mM sodium acetate, pH 4.5. The solution was then applied on a Q sepharose column, previously equilibrated in 10 mM Piperazin, Sigma, pH 5.5. Unbound material is washed out with the equilibration buffer. Bound proteins are eluted with a linear gradient of 0-0.3 M Sodium chloride in 10 mM Piperazin, pH 5.5 over 10 column volumes. Glucoamylase containing fractions are collected and the purity was confirmed by SDS-PAGE.

T_{1/2} (half-life) Method I

The thermal stability of variants is determined as T_{1/2} using the following method: 950 microliter 50 mM sodium acetate buffer (pH 4.3) (NaOAc) is incubated for 5 minutes at 68°C or 70°C. 50 microliter enzyme in buffer (4 AGU/ml) is added. 2 x 40 microliter samples are taken at 0, 5, 10, 20, 30 and 40 minutes and chilled on ice. The activity (AGU/ml) measured before incubation (0 minutes) is used as reference (100%). The decline in stability (in percent) is calculated as a function of the incubation time. The % residual glucoamylase activity is determined at different times. T_{1/2} is the period of time until which the % relative activity is decreased to 50%.

T_{1/2} (half-life) (Method II)

The T_{1/2} is measured by incubating the enzyme (ca 0.2 AGU/ml) in question in 30% glucose, 50 mM Sodium acetate at pH 4.5 at the temperature in question (e.g., 70°C). Samples are withdrawn at set time intervals and chilled on ice and residual enzyme activity measured by the pNPG method (as described below).

The % residual glucoamylase activity is determined at different times. T_{1/2} is the period of time until which the % relative activity is decreased to 50%.

Residual Enzyme Activity (pNPG method)

pNPG reagent:

0.2 g pNPG (p-nitrophenylglucopyranoside) is dissolved in 0.1 M acetate buffer (pH 4.3) and made up to 100 ml.

Borate solution:

3.8g Na₂B₄O₇ · 10 H₂O is dissolved in Milli-Q water and made up to 100 ml.

25 microL samples are added 50 microL substrate and incubated 2 hr at 50°C. The reaction is stopped by adding 150 microL ml borate solution. The optical density is measured at 405 nm, and the residual activity calculated.

Construction Of pAMGY

The pAMGY vector was constructed as follows: The lipase gene in pJSO026 was replaced by the AMG gene, which was PCR amplified with the forward primer; FG2: 5'-CAT CCC CAG GAT CCT TAC TCA GCA ATG-3' and the reverse primer: RG2: 5'-CTC AAA CGA CTC ACC AGC CTC TAG AGT-3' using the template plasmid pLAC103 containing the AMG gene. The pJSO026 plasmid was digested with XbaI and SmaI at 37°C for 2 hours and the PCR amplicon was blunt ended using the Klenow fragment and then digested with XbaI. The vector fragment and the PCR amplicon were ligated and transformed into *E.coli* by electrotransformation. The resulting vector is designated pAMGY.

Construction Of pLaC103

The *A. niger* AMGII cDNA clone (Boel et al., (1984), supra) is used as source for the construction of pLaC103 aimed at *S. cerevisiae* expression of the GII form of AMG.

The construction takes place in several steps, outlined below.

pT7-212 (EP37856/ US patent no. 5162498) is cleaved with XbaI, blunt-ended with Klenow DNA polymerase and dNTP. After cleavage with EcoRI the resulting vector fragment is purified from an agarose gel-electrophoresis and ligated with the 2.05 kb EcoRI-EcoRV fragment of pBoel53, thereby recreating the XbaI site in the EcoRV end of the AMG encoding fragment in the resulting plasmid pG2x.

In order to remove DNA upstream of the AMG cds, and furnish the AMG encoding DNA with an appropriate restriction endonuclease recognition site, the following construct was made:

The 930 bp EcoRI-PstI fragment of p53 was isolated and subjected to AluI cleavage, the resulting 771 bp Alu-PstI fragment was ligated into pBR322 with blunt-ended EcoRI site (see above) and cleaved with PstI. In the resulting plasmid pBR-AMG', the EcoRI site was recreated just 34 bp from the initiation codon of the AMG cds.

From pBR-AMG' the 775 bp EcoRI - PstI fragment was isolated and joined with the 1151 bp PstI - XbaI fragment from pG2x in a ligation reaction including the XbaI - EcoRI vector fragment of pT7-212.

The resulting plasmid pT7GII was submitted to a BamHI cleavage in presence of alkaline phosphatase followed by partial SphI cleavage after inactivation of the phosphatase. From this reaction was the 2489 bp SphI-BamHI fragment, encompassing the S.c. TPI promoter linked to the AMGII cds.

The above fragment together with the 1052 bp BamHI fragment of pT7GII was ligated with the alkaline phosphatase treated vector fragment of pMT743 (EP37856/US 5162498), resulting from SphI-BamHI digestion. The resulting plasmid is pLaC103.

5 Screening For Thermostable AMG Variants

The libraries are screened in the thermostable filter assay described below.

Filter Assay For Thermostability

Yeast libraries are plated on a sandwich of cellulose acetate (OE 67, Schleicher & Schuell, Dassel, Germany) - and nitrocellulose filters (Protran-Ba 85, Schleicher & Schuell, Dassel, Germany) on SCFura⁺ agar plates with 100 µg/ml ampicillin at 30°C for at least 72 hrs. The colonies are replica plated to PVDF filters (Immobilon-P, Millipore, Bedford) activated with methanol for 1 min or alternatively a Protran filter (no activation) and subsequently washed in 0.1 M NaAc and then incubated at room temperature for 2 hours. Colonies are washed from PVDF/Protran filters with tap water. Each filter sandwiches and PVDF/Protran filters are specifically marked with a needle before incubation in order to be able to localise positive variants on the filters after the screening. The PVDF filters with bound variants are transferred to a container with 0.1 M NaAc, pH 4.5 and incubated at 47°C or alternatively 67-69°C in case of Protran filters for 15 minutes. The sandwich of cellulose acetate and nitrocellulose filters on SC ura-agar plates are stored at room temperature until use. After incubation, the residual activities are detected on plates containing 5% maltose, 1% agarose, 50 mM NaAc, pH 4.5. The assay plates with PVDF filters are marked the same way as the filter sandwiches and incubated for 2 hrs. at 50°C. After removal of the PVDF filters, the assay plates are stained with Glucose GOD perid (Boehringer Mannheim GmbH, Germany). Variants with residual activity are detected on assay plates as dark green spots on white background. The improved variants are located on the storage plates. Improved variants are rescreened twice under the same conditions as the first screen.

30 General Method For Random Mutagenesis By Use Of The DOPE Program

The random mutagenesis may be carried out by the following steps:

1. Select regions of interest for modification in the parent enzyme,
2. Decide on mutation sites and non-mutated sites in the selected region,
3. Decide on which kind of mutations should be carried out, *e.g.*, with respect to the desired stability and/or performance of the variant to be constructed,
4. Select structurally reasonable mutations,

5. Adjust the residues selected by step 3 with regard to step 4.
6. Analyze by use of a suitable dope algorithm the nucleotide distribution.
7. If necessary, adjust the wanted residues to genetic code realism, *e.g.* taking into account constraints resulting from the genetic code, *e.g.* in order to avoid introduction of stop codons; the skilled person will be aware that some codon combinations cannot be used in practice and will need to be adapted
8. Make primers
9. Perform random mutagenesis by use of the primers
10. Select resulting glucoamylase variants by screening for the desired improved properties.

Dope Algorithm

Suitable dope algorithms for use in step 6 are well known in the art. One such algorithm is described by Tomandl, D. et al., 1997, Journal of Computer-Aided Molecular Design 11:29-38. Another algorithm is DOPE (Jensen, LJ, Andersen, KV, Svendsen, A, and Kretzschmar, T (1998) Nucleic Acids Research 26:697-702).

Method Of Extracting Important Regions For Temperature Activity Using Molecular Simulation.

The X-ray structure and/or the model-build structure of the enzyme of interest, here AMG, are subjected to molecular dynamics simulations. The molecular dynamics simulation are made using the CHARMM (from Molecular simulations (MSI)) program or other suitable programs, e.g., DISCOVER (from MSI). The dynamics are made in vacuum, or including crystal waters, or with the enzyme in question embedded in a suitable waters, e.g., a sphere or a box. The simulation are run for 300 picoseconds (ps) or more, e.g., 300-1200 ps. The isotropic fluctuations are extracted for the CA carbons of the structures and comparison between the structures are made. More details on how to get the isotropic fluctuations can be found in the CHARMM manual (available from MSI) and hereby incorporated herein by reference.

The molecular dynamics simulation can be carried out using standard charges on the chargeable amino acids. For instance, Asp and Glu is negatively charged and Lys and Arg are positively charged. This condition resembles the medium pH of approximately 7.0. To analyze a lower pH, titration of the molecule can be done to obtain the altered pKa's of the normal titrateable residues within pH 2-10; Lys, Arg, Asp, Glu, Tyr and His. Also Ser, Thr and Cys are titrateable but are not taking into account here. Here the altered charges due to the pH has been described as all Arg, Lys negative at high pH, and all Asp, Glu are uncharged. This imitates a pH around 4 to 5 where the titration Asp and Glu normally takes place.

Model building of the enzyme of interest can be obtained by using the HOMOLOGY model in the MSI program package. The crystal structure of *Aspergillus awamori* variant X100 can be found in, e.g., 3GLY and 1DOG in the Brookhaven database.

EXAMPLES

EXAMPLE 1

Construction of AMG G2 variants

5 Site-directed mutagenesis:

For the construction of variants of AMG G2 (SEQ ID NO: 2) the commercial kit, Chameleon double-stranded, site-directed mutagenesis kit was used according to the manufacturer's instructions.

10 The gene encoding the AMG G2 enzyme in question is located on pMT838 prepared by deleting the DNA between G2 nt. 1362 and G2 nt. 1530 in plasmid pCAMG91 (see Figure 1) comprising the AMG G1 form.

In accordance with the manufacturer's instructions the Scal site of the Ampicillin gene of pMT838 was changed to a MluI site by use of the following primer:

7258: 5'p gaa tga ctt ggt tga cgc gtc acc agt cac 3' (SEQ ID NO: 3).

15 (Thus changing the Scal site found in the ampicillin resistance gene and used for cutting to a MluI site). The pMT838 vector comprising the AMG gene in question was then used as a template for DNA polymerase and oligo 7258 (SEQ ID NO: 3) and 21401 (SEQ ID NO: 4).

Primer no. 21401 (SEQ ID NO: 4) was used as the selection primer.

20 21401: 5'p gg gga tca tga tag gac tag cca tat taa tga agg gca tat acc acg cct tgg acc tgc gtt ata gcc 3'

(Changes the Scal site found in the AMG gene without changing the amino acid sequence).

The desired mutation (e.g., the introduction of a cystein residue) is introduced into the
25 AMG gene in question by addition of appropriate oligos comprising the desired mutation.

The primer 107581 was used to introduce T12P

107581: 5` pgc aac gaa gcg ccc gtg gct cgt ac 3` (SEQ ID NO: 5)

30 The mutations are verified by sequencing the whole gene. The plasmid was transformed into *A. oryzae* using the method described above in the "Materials & Methods" section. The variant was fermented and purified as described above in the "Materials & Methods" section.

EXAMPLE 2

Construction, by localized random, doped mutagenesis, of *A. niger* AMG variants having improved thermostability compared to the parent enzyme

- 5 To improve the thermostability of the *A. niger* AMG random mutagenesis in pre-selected region was performed.

Residue:

Region: L19-G35

Region: A353-V374

- 10 The DOPE software (see Materials and Methods) was used to determine spiked codons for each suggested change in the above regions minimizing the amount of stop codons (see table 1). The exact distribution of nucleotides was calculated in the three positions of the codon to give the suggested population of amino acid changes. The doped regions were doped specifically in the indicated positions to have a high chance
15 of getting the desired residues, but still allow other possibilities.

The first column is the amino acid to be mutated, the second column is the percentage of wild type and the third column defined the new amino acid(s).

Table 1

20 Doping in L19-G35

	L19	90%	N
	N20	95%	T
	N21	Constant	
	I22	Constant	
25	G23	95%	A
	A24	90%	S,T
	D25	93%	S,T,R
	G26	95%	A
	A27	90%	S,T
30	W28	<80%	R,Y
	V29	Constant	
	S30	93%	T,N
	G31	95%	A
	A32	95%	V
35	D33	80%	R,K,H
	S34	90%	N

G35 Constant

The resulting doped oligonucleotide strand is shown in table 2 as sense strand:
with the primer sequence, the wild type nucleotide sequence, the parent amino acid
5 sequence and the distribution of nucleotides for each doped position.

Table 2:

	Position:	19	20	21	22	23	24	25	26	27
	A.a. seq.:	L	N	N	I	G	A	D	G	A
10	primer:	12T	A3T	AAC	ATC	G4G	5CG	67C	G4T	8CT
	wt. seq.:	CTG	AAT	AAC	ATC	GGG	GCG	GAC	GGT	GCT
	Pos. (cont.):	28		29	30	31	32	33	34	35
	A.a. (cont.):	W	V	S	G	A		D	S	G
15	primer:		91010	GTG	1112C	G4C	G13G	141516	1718T	GGC
	Wt seq.:		TGG	GTG	TCG	GGC	GCG	GAC	TCT	GGC

Distribution of nucleotides for each doped position.

- 1: A10,C90
- 20 2: A6, T94
- 3: A95,C5
- 4:G95,C5
- 5:G91,A3,T3,C3
- 6:G95,A3,C2
- 25 7:G3,A95,C2
- 8:G92,A4,T4
- 9:A3,T97
- 10:G95,T5
- 11:G3,A97
- 30 12:G95,A2,C3
- 13:T5,C95
- 14:G88,A8,C4
- 15:G7,A93
- 16:G4,C96
- 35 17:G4,A96
- 18:G95,A2,C3

Forward primer (SEQ ID NO: 6):

FAMGII '5-C GAA GCG ACC GTG GCT CGT ACT GCC ATC 12T A3T AAC ATC G4G
5CG 67C G4T 8CT 91010 GTG 1112C G4C G13G 141516 1718T GGC ATT GTC GTT

5 GCT AGT CCC AGC ACG GAT AAC-3'

Reverse primer (SEQ ID NO: 7):

RAMG1: 5'-GAT GGC AGT ACG AGC CAC GGT CGC TTC G-3'

10 Table 3_

Doping in region A353-V374:

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30

A353	<80%	D,E,Q,N,Y
L354	90%	Q,E
Y355	90%	N,Q
15 S356	90%	T,D,N
G357	80%	P,A,S,T
A358	93%	S
A359	90%	S,T,N
T360	90%	R,K
20 G361	85%	A,S,T
T362	90%	S
Y363	Constant	
S364	93%	D
S365	93%	N,Q,K
25 S366	93%	P,D
S367	Constant	
S368	93%	D,N,T
T369	93%	Q,E
Y370	Constant	
30 S371	93%	N
S372	93%	N,T
I373	Constant	
V374	93%	N,Y,H

The resulting doped oligonucleotide strand is shown in table 4 as sense strand:
with the primer sequence, wild type nucleotide sequence, the parent amino acid
sequence and the distribution of nucleotides for each doped position.

5 Table 4:

	Position:	353	354	355	356	357	358	359	360	361	362
	A.a. seq.:	A	L	Y	S		D	A	A	T	G
	primer:	123	45A	6AC	78C	910T	11CT	1213T	1415A	1617C	18CC
	Wt. seq.:	GCA	CTG	TAC	AGC	GAT	GCT	GCT	ACT	GGC	ACC
10	Pos. (cont.):		363		364		365		366	367368	369
	A.a. seq. (cont.):	Y		S		S		S		T	Y
	primer (cont.):		TAC	1920T	A2122	2324C	AGT	1425C	2627G	T28T	
	wt. Seq.(cont.):		TAC	TCT	TCG	TCC	AGT	TCG	ACT	TAT	
	Pos. (cont.):		371		372		373	374			
15	A.a. pos. (cont.):	S		S		I	V				
	primer (cont.)			A16T	2930T	ATT	313233				
	wt. Seq.(cont.):		AGT	AGC	ATT	GTA					

Distribution of nucleotides for each doped position.

- 20 1:G91,A3,T3,C3
- 2:A13,C87
- 3:A40,T60
- 4:G3,A3,C94
- 5:A6,T94
- 25 6:G4,A4,T92
- 7:G2,A96,C2
- 8:G93,A3.5,C3.5
- 9:G87,A8,C5
- 10:A84,C16
- 30 11:G93,T7
- 12:G92,A5,T3
- 13:A3,C97
- 14:G3,A97
- 15:G2,A2,T4,C92
- 35 16:G93,A7
- 17:G93,C7

18:A90,T10
 19:G4,A96
 20:G95,A5
 21:G96,A4
 5 22:G3,C97
 23:G2,A1,T95,C2
 24:A3,C97
 25:G95,A3,C2
 26:G2,A96,C2
 10 27:A5,C95
 28:A95,T5
 29:G2,A98
 30:G94,A4,C2
 31:G94,A3,T1,C2
 15 32:A4,T96
 33:A20,C80

Primer: FAMGIV (SEQ ID NO: 8)
 5'-GTG TCG CTG GAC TTC TTC AAG 123 45A 6AC 78C 910T 11CT 1213T 1415A
 20 1617C 18CC TAC 1920T A2122 2324C AGT 1425C 2627G T28T A16T 2930C ATT
 313233 GAT GCC GTG AAG ACT TTC GCC GA-3'

Primer RAMGVI (SEQ ID NO: 9)
 5'-ctt gaa gaa gtc cag cga cac-3'

Random mutagenesis

The spiked oligonucleotides apparent from Table 2 and 3 (which by a common term is designated FAMG) and reverse primers RAMG for the L19-G35 region and specific SEQ ID NO: 2 primers covering the N-terminal (FG2: 5'- CAT CCC CAG GAT CCT TAC
 30 TCA GCA ATG-3' (SEQ ID NO: 10) and C-terminal (RG2: 5'- CTC AAA CGA CTC ACC
 AGC CTC TAG AGT (SEQ ID NO: 11) are used to generate PCR-library-fragments by the overlap extension method (Horton et al., Gene, 77 (1989), pp. 61-68) with an overlap of 21 base pairs. Plasmid pAMGY is template for the Polymerase Chain Reaction. The PCR fragments are cloned by homologous recombination in the *E. coli*/yeast shuttle vector
 35 pAMGY (see Materials and Methods).

Screening

The library was screened in the thermostability filter assays using a Protran filter and incubating at 67-69°C as described in the "Material & Methods" section above

5

EXAMPLE 3

Construction, by PCR shuffling spiked with DNA oligos, of A. niger AMG variants having improved thermostability compared to the parent enzyme.

The polymerase chain reaction (PCR) method was used to prepare DNA
10 fragments carrying the AMG gene and flanking regions. Approximately 10 ug DNA was digested with Dnase, and run on a 2 % agarose gel. Fragments of 50-150 bp were purified from the gel. Approximately 1 ug purified fragments were mixed with a 5-15 fold molar excess of oligos carrying the desired mutations. The oligos were of the following kind (for the construction of Hklib1, Hklib2, Hklib3 etc., respectively):

10
15
20
25
30
35

Hklib1:

Hk1-T2X:

5'-ATGTGATTTCCAAGCGCGCGVNNNTTGGATTCATGGTTGAGCAA (SEQ ID NO:14)

Hk1-N9X:

5'-CCTTGGATTCATGGTTGAGCVNNGAAGCGACCGTGGCTCGTAC (SEQ ID NO:15)

Hk1-A11X:

25 5'-ATTCATGGTTGAGCAACGAADVNNACCGTGGCTCGTACTGCCAT (SEQ ID NO:16)

Hk1-L66X:

5'-TCCTCAAGACCCTCGTCGATVNNTTCCGAAATGGAGATACCAG (SEQ ID NO:17)

Hk1-S386X:

30 5'-CTTTGCGCGATGGCTTCGTCVNNATTGTGGAACTCACGCCGC (SEQ ID NO:18)

Hk1-E389X:

35 5'-ATGGCTTCGTCTCTATTGTGVNNACTCACGCCGCAAGCAACGG (SEQ ID NO:19)

Hk1-T390X:

5'-GCTTCGTCTCTATTGTGGA**VNN**CACGCCGCAAGCAACGGCTC (SEQ ID NO:20)

Hk1-A393X:

5'-CTATTGTGGA**AACT**CACGCC**VNN**AGCAACGGCTCCATGTCCGA (SEQ ID NO:21)

Hk1-S394X:

5'-TTGTGGA**AACT**CACGCCGCA**VNNA**ACGGCTCCATGTCCGAGCA (SEQ ID NO:22)

Hk1-N395X:

10 5'-TGGA**AACT**CACGCCGCAAGC**VNNG**GCTCCATGTCCGAGCAATA (SEQ ID NO:23)

Hk1-G396X:

5'-**AACT**CACGCCGCAAGCAAC**VNNT**CCATGTCCGAGCAATACGA (SEQ ID NO:24)

Hk1-K404X:

5'-CCATGTCCGAGCAATACGAC**VNNT**CTGATGGCGAGCAGCTTTC (SEQ ID NO:25)

Hk1-D406X:

5'-CCGAGCAATACGACAAGTCT**VNNG**GCGAGCAGCTTTCCGCTCG (SEQ ID NO:26)

Hk1-E408X:

5'-AATACGACAAGTCTGATGGC**VNN**CAGCTTTCCGCTCGCGACCT (SEQ ID NO:27)

Hk1-L410X:

5'-ACAAGTCTGATGGCGAGCAG**VNNT**CCGCTCGCGACCTGACCT (SEQ ID NO:28)

Hk1-L423X:

5'-CCTGGTCTTATGCTGCTCTG**VNN**ACCGCCAACAACCGTCGTAA (SEQ ID NO:29)

Hk1-N426X:

5'-ATGCTGCTCTGCTGACCGCC**VNNA**ACCGTCGTAACTCCGTCGTG (SEQ ID NO:30)

Hk1-N427X:

5'-CTGCTCTGCTGACCGCCAAC**VNN**CGTCGTAACTCCGTCGTGCCT (SEQ ID NO:31)

Hk1-Y402X:

5'-ACGGCTCCATGTCCGAGCAANN**C**GACAAGTCTGATGGCGAGCAGCT (SEQ ID NO:32)

5 **Hklib2:**

Hk2-L234X-SENSE:

5'-CTGGACCGGCAGCTTCATT**NNK**GCCAACTTCGATAGCAGCC (SEQ ID NO:33)

Hk2-A235S-ANTISENSE:

10 5'-GAACGGCTGCTATCGAAGTT**AGAC**AGAATGAAGCTGCCGGTC (SEQ ID NO:34)

Hk2-F237X-SENSE:

5-CAGCTTCATTCTGGCCAAC**NAT**GATAGCAGCCGTTCCGGCA (SEQ ID NO:35)

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165

Hk2-T310S-ANTISENSE:

5'-ACGGGTTGCCGTTGTAGTA**A**AGTCCTCAGGGTACCGACCC (SEQ ID NO:44)

Hk2-Y311N-SENSE:

5 5'-TCGGTACCCTGAGGACACGA**A**ATTACAACGGCAACCCGTGGT (SEQ ID NO:45)

Hk2-Y312Q-ANTISENSE:

5'-GGAACCACGGGTTGCCGTT**T**TGGTACGTGTCCTCAGGGTAC (SEQ ID NO:46)

10 **Hk2-Y312N-ANTISENSE:**

5'-GGAACCACGGGTTGCCGTT**A**TTGTACGTGTCCTCAGGGTAC (SEQ ID NO:47)

Hk2-N313T-SENSE:

5'-CCCTGAGGACACGTACTAC**A**CTGGCAACCCGTGGTTCCTGT (SEQ ID NO:48)

Hk2-N313S-SENSE:

5'-CCCTGAGGACACGTACTACT**C**TGGCAACCCGTGGTTCCTGT (SEQ ID NO:49)

Hk2-N313G-SENSE:

5'-CCCTGAGGACACGTACTAC**G**GTGGCAACCCGTGGTTCCTGT (SEQ ID NO:50)

Hk2-N315Q-ANTISENSE:

5'-AGGTGCACAGGAACCACGGTT**G**GCCGTTGTAGTACGTGTCC (SEQ ID NO:51)

25 **Hk2-N315E-ANTISENSE:**

5'-AGGTGCACAGGAACCACGGTT**C**GCCGTTGTAGTACGTGTCC (SEQ ID NO:52)

Hk2-N315R-ANTISENSE:

5'-AGGTGCACAGGAACCACGGT**C**TGCCGTTGTAGTACGTGTCC (SEQ ID NO:53)

Hk2-F318Y-ANTISENSE:

5'-CGGCAGCCAAGGTGCACAG**A**TACCACGGGTTGCCGTTGTAG (SEQ ID NO:54)

Hk2-Q409P-SENSE:

35 5'-CGACAAGTCTGATGGCGAG**C**CACTTTCCGCTCGCGACCTGA (SEQ ID NO:55)

Hklib3:

Hk3-D336X-SENSE:

5'-CGATGCTCTATACCAAGTGG**NN**KAAGCAGGGGTCGTTGGAGG (SEQ ID NO:56)

Hk3-K337X-SENSE:

5'-TGCTCTATACCAAGTGGGAC**NN**KCAGGGGTCGTTGGAGGTCA (SEQ ID NO:57)

Hk3-Q338X-ANTISENSE:

5'-CTGTGACCTCCAACGACCC**G**NNCTTGTCCCACTGGTATAGA (SEQ ID NO:58)

Hk3-G339X-SENSE:

10 5'-ATACCAAGTGGGACAAGCAG**N**CUTC GTTGGAGGTCACAGATG (SEQ ID NO:59)

Hk3-S340X'-ANTISENSE:

5'-ACACATCTGTGACCTCCAA**A**NTCCCCTGCTTGTCCCACTGG (SEQ ID NO:60)

Hk3-S340X''-ANTISENSE:

5'-ACACATCTGTGACCTCCAA**A**NCCCCCTGCTTGTCCCACTGG (SEQ ID NO:61)

Hk3-L341X-SENSE:

5'-GTGGGACAAGCAGGGGTCG**NU**UGAGGTCACAGATGTGTCGC (SEQ ID NO:62)

Hk3-K352Q-SENSE:

5'-TGTGTCGCTGGACTTCTT**CA**AGCACTGTACAGCGATGCTG (SEQ ID NO:63)

Hk3-K352R-SENSE:

25 5'-TGTGTCGCTGGACTTCTT**CA**GAGCACTGTACAGCGATGCTG (SEQ ID NO:64)

Hk3-A353D-ANTISENSE:

5'-TAGCAGCATCGCTGTACAG**A**TCCTTGAAGAAGTCCAGCGAC (SEQ ID NO:65)

Hk3-A353S-ANTISENSE:

30 5'-TAGCAGCATCGCTGTACAG**A**GACTTGAAGAAGTCCAGCGAC (SEQ ID NO:66)

Hk3-S356P-SENSE:

5'-ACTTCTTCAAGGCACTGTAC**CC**AGATGCTGCTACTGGCACCT (SEQ ID NO:67)

Hk3-S356N-SENSE:

5'-ACTTCTTCAAGGCACTGTAC**AA**UGATGCTGCTACTGGCACCTA (SEQ ID NO:68)

Hk3-S356D-SENSE:

5'-ACTTCTTCAAGGCACTGTAC**GAUG**ATGCTGCTACTGGCACCTA (SEQ ID NO:69)

Hk3-D357S-ANTISENSE:

5 5'-GAGTAGGTGCCAGTAGCAGC**AGAG**CTGTACAGTGCCTTGAAGA (SEQ ID NO:70)

Hk3-A359S-SENSE:

5'-GGCACTGTACAGCGATGCTT**CT**ACTGGCACCTACTCTTCGT (SEQ ID NO:71)

10

Hk3-T360V-ANTISENSE:

5'-TGGACGAAGAGTAGGTGCC**AAC**AGCAGCATCGCTGTACAGT (SEQ ID NO:72)

Hk3-G361X-SENSE:

15 5'-TGTACAGCGATGCTGCTACT**NCT**ACCTACTCTTCGTCCAGTTC (SEQ ID NO:73)

Hk3-T362R-ANTISENSE:

5'-GTCGAACTGGACGAAGAGTAT**CT**GCCAGTAGCAGCATCGCTG (SEQ ID NO:74)

20 **Hk3-S364X-SENSE:**

5'-TGCTGCTACTGGCACCTAC**NNK**TCGTCCAGTTCGACTTATAG (SEQ ID NO:75)

Hk3-S365X-SENSE:

5'-TGCTACTGGCACCTACTCT**NNK**TCCAGTTCGACTTATAGTAG (SEQ ID NO:76)

25

Hk3-S366T-ANTISENSE:

5'-ATGCTACTATAAGTCGAACT**AG**TCGAAGAGTAGGTGCCAGTA (SEQ ID NO:77)

Hk3-S368X-ANTISENSE:

30 5'-TCTACAATGCTACTATAAGT**AGN**ACTGGACGAAGAGTAGGTG (SEQ ID NO:78)

Hk3-T369X-SENSE:

5'-CTACTCTTCGTCCAGTTCG**NNK**TATAGTAGCATTGTAGATGCC (SEQ ID NO:79)

35 **Hk3-S371X-ANTISENSE:**

5'-TTCACGGCATCTACAATGCT**ATN**ATAAGTCGAACTGGACGAAG (SEQ ID NO:80)

Hk3-S372X-SENSE:

5'-CGTCCAGTTCGACTTATAGT**NNT**ATTGTAGATGCCGTGAAGAC (SEQ ID NO:81)

- 5 To the mix of Dnase treated DNA and oligos was added nucleotides, PCR buffer and Taq/Pwo polymerase. A PCR assembly reaction was performed, using first 94 °C for 2 min., then 35-40 cycles with the following incubation times: 94°C, 30 sec.; 45 °C, 30 sec.; 72°C, 60 sec ; then finally 72°C for 5 min.

10 An PCR amplification reaction was performed with 1 uL of the assembly reaction as template, and adding primers that anneal to the regions flanking the AMG gene.

Parameters: first 94 °C for 2 min., then 35-40 cycles with the following incubation times: 94°C, 30 sec.; 55°C, 30 sec.; 72 °C, 90 sec ; then finally 72°C for 10 min.

The resulting PCR product was purified from a 1% agarose gel, mixed with linearized vector and transformed into competent yeast cells, as described above.

EXAMPLE 4

Specific Activity

15 AMG G2 variants were constructed as described above in Example 1. The specific activity as k_{cat} were measured on purified samples at pH 4.3, 37°C, using maltose and
20 maltoheptaose as substrate as described in the "Materials & Methods" section above. The specific activity as AGU/mg were also measured at pH 4.3, 37°C, using maltose as substrate as described in the "Materials & Methods" section above.

Variant	Kcat (sec.-1)	
	Maltose	Maltoheptaose
AMG G2 (wt)	6.0	38
N110T	9.7	27.8
V111P	12.0	43.2
S119P	6.2	44.0
G127A	21.0	40.0
G207N	30.5	36.3

Variant	AGU/mg
---------	--------

AMG G2 (wild type)	1.8
N110T	3.5
V111P	3.1
S119P	2.1
G127A	5.8
G207N	5.7
L3N	2.3
S56A	2.6
A102*	2.5
D403S	2.2
I18V+T51S+S56A+V59T+L60A	3.3
S119P+Y402F	2.7
S119P+I189T+Y223F+F227Y+Y402F	3.0

EXAMPLE 5

Thermostability at 70°C

An AMG G2 S119P variant was constructed using the approach described in Example 1.

The thermostability was determined as $T_{1/2}$ using Method I, and as % residual activity after incubation for 30 minutes in 50 mM NaOAc, pH 4.5, 70°C, 0.2 AGU/ml, as described in the "Material & Methods" section above. The result of the tests are listed in the Table below and compared to the wild-type *A. niger* AMG G2.

<i>A. niger</i> AMG (Enzyme)	Residual activity (%)	$T_{1/2}$ (min.)
S119P variant	22	17
wild-type (SEQ ID NO: 2)	13	8

EXAMPLE 6

Thermostability at 68°C

AMG G2 variants were constructed using the approach described in Example 3, except for variants nos. 1 and 2 in the Table below, which were prepared by shuffling as described in WO 95/22625 (from Affymax Technologies N.V.).

The thermostability was determined as $T_{1/2}$ using method I at 68°C as described in the "Materials & Methods" section and compared to the wild-type *A. niger* AMG G2 under

the same conditions. Evaluation of variants were performed on culture broth after filtration of the supernatants.

	Variant	T _{1/2} (min)	T _{1/2} <i>A. niger</i> AMG G2 (wild type) (min)
1	A246T+T72I	11.3	8.5
2	G447S+S119P	11.4	7.9
3	E408R+A425T+S465P+T494A	8.6	8.1
4	E408R+S386N	12.6	8.9
5	T2P	9.3	8.5
6	T2Q+A11P+S394R	10.7	8.5
7	T2H	9.5	8.9
8	A11E+E408R	12.7	9.3
9	T2M+N9A+T390R+D406N+L410R	10.7	8.5
10	A393R	17.7	8.4
11	T2R+S386R+A393R	14.1	8.4
12	A393R+L410R	14.7	7.9
13	A1V+L66R+Y402F+N427S+S486G	11.7	8.5
14	T2K+S30P+N427M+S444G+V470 M	11.4	8.4

5

Thermostability at 70°C on purified samples.

	Enzyme	T _{1/2} (min)
15	AMG G2 (wild type)	7.4
16	T2E+T379A+S386K+A393R	11.6
17	E408R+S386N	10.2
18	T2Q+A11P+S394R	9.8
19	A1V+L66R+Y402F+N427S+S486G	14.1
20	A393R	14.6
21	T2R+S386R+A393R	14.1
22	A393R+L410R	12.9
23	Y402F	10.1

EXAMPLE 7

Thermostability at 68°C

AMG G2 variants were constructed by shuffling using the approach described in Example 3 followed by shuffling of positive variants.

- 5 The thermostability was determined as $T_{1/2}$ using method I at 68°C as described in the “Materials & Methods” section and compared to the wild-type *A. niger* AMG G2 under the same conditions. Evaluation of variants were performed on culture broth after filtration of the supernatants.

	Variant	$T_{1/2}$ (min)	$T_{1/2}$ <i>A. niger</i> AMG G2 (wild type) (min)
24	PLASD ⁱ +V59A+A393R+T490A	27.2	6.8

10 i = N-terminal extension

EXAMPLE 8

Thermostability at 68°C

- 15 AMG G2 variants were constructed using the approach described in Example 3. The thermostability was determined as $T_{1/2}$ using method I at 68°C as described in the “Materials & Methods” section and compared to the wild-type *A. niger* AMG G2 under the same conditions. Evaluation of variants were performed on culture broth after filtration of the supernatants.

	Variant	$T_{1/2}$ (min)	$T_{1/2}$ <i>A. niger</i> AMG G2 wild-type (min)
25	D357S+T360V+S371H	6.6	5.9
26	N313G+F318Y	8.9	5.9
27	S356P+S366T	7.3	5.8
28	S340G+D357S+T360V+S386P	7.2	5.8

EXAMPLE 9

Thermostability at 70°C

An AMG G2 variants was constructed using the approach described in Example 1 and evaluated as semi-purified (filtration of culture broth followed by desalting on a G-25 column) samples.

The thermostability was determined as % residual activity using Method I in 50 mM NaOAc, pH 4.5, 70°C, as described in the "Material & Methods" section above. The result of the test is listed in the Table below and compared to the wild-type *A. niger* AMG G2.

	Enzyme	T½ (min)
29	AMG G2 (wild type)	7
30	Y402F+S411V	60
31	S119P+Y402F+S411V	115
32	S119P+Y312Q+Y402F+T416H	50

EXAMPLE 10

Thermostability at 70°C in presence of 30% glucose

AMG G2 variants were constructed using the approach described in Example 3.

The thermostability was determined as T½ using method II at 70°C as described in the "Materials & Methods" section and compared to the wild-type *A. niger* AMG G2 under the same conditions.

	Enzyme	T½ (hr)
33	AMG G2 (wild type)	1.5
34	Y402F	2.5
35	A393R	4.0
36	T2R+S386R+A393R	2.0
37	PLASD(N-terminal)+V59A+A393R+T490A	16.0

EXAMPLE 11

Saccharification performance of AMG variants S119P+Y402F+S411V and PLASD(N-terminal)+V59A+A393R+T490A, respectively.

Saccharification performance of the AMG variants S119P+Y402F+S411V and PLASD(N-terminal)+V59A+A393R+T490A, respectively, both having improved thermostability are tested at 70°C as described below.

Reference enzyme is the wild-type *A. niger* AMG G2. Saccharification is run under the following conditions:

Substrate	10 DE Maltodextrin, approx. 30% DS (w/w)
Temperature	70°C
Initial pH	4.3 (at 70°C)
Enzyme dosage	0.24 AGU/g DS

Saccharification

The substrate for saccharification is made by dissolving maltodextrin (prepared from common corn) in boiling Milli-Q water and adjusting the dry substance to approximately 30% (w/w). pH is adjusted to 4.3. Aliquots of substrate corresponding to 15 g dry solids are transferred to 50 ml blue cap glass flasks and placed in a water bath with stirring. Enzymes are added and pH re-adjusted if necessary. The experiment is run in duplicate. Samples are taken periodically and analysed at HPLC for determination of the carbohydrate composition.